

KD-Validated Anti-EHMT2/G9A Rabbit Monoclonal Antibody

Rabbit monoclonal antibody
Catalog # AGI2359

Product Information

Application	WB, FC
Primary Accession	Q96KQ7
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	23GB820
Calculated MW	132370
Gene Name	EHMT2
Aliases	KMT1C; G9A; C6orf3; BAT8; Euchromatic Histone-Lysine N-Methyltransferase; Histone-Lysine N-Methyltransferase EHMT2; Histone H3-K9 Methyltransferase; HLA-B Associated Transcript 8; Lysine N-Methyltransferase 1C; H3-K9-HMTase; Em:AF134726.3; NG36/G9a; NG36; Histone-Lysine N-Methyltransferase, H3 Lysine-9 Specific 3; Chromosome 6 Open Reading Frame 30; Ankyrin Repeat-Containing Protein; G9A Histone Methyltransferase; HLA-B-Associated Transcript 8; EC 2.1.1.367; Protein G9a; EC 2.1.1.-; GAT8
Immunogen	A synthesized peptide derived from human EHMT2/G9A

Additional Information

Gene ID	10919
Other Names	Histone-lysine N-methyltransferase EHMT2, 2.1.1.-, 2.1.1.367, Euchromatic histone-lysine N-methyltransferase 2, HLA-B-associated transcript 8, Histone H3-K9 methyltransferase 3, H3-K9-HMTase 3, Lysine N-methyltransferase 1C, Protein G9a, EHMT2, BAT8, C6orf30, G9A, KMT1C, NG36

Protein Information

Name	EHMT2
Synonyms	BAT8, C6orf30, G9A, KMT1C, NG36
Function	Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones (PubMed: 11316813 , PubMed: 20084102). Also mediates monomethylation of 'Lys-56' of histone H3 (H3K56me1) in G1 phase, leading to promote interaction between histone H3 and PCNA and regulating DNA replication (PubMed: 22387026). Also weakly methylates 'Lys-27' of histone H3 (H3K27me) (PubMed: 11316813). Also

required for DNA methylation, the histone methyltransferase activity is not required for DNA methylation, suggesting that these 2 activities function independently. Probably targeted to histone H3 by different DNA-binding proteins like E2F6, MGA, MAX and/or DP1. Also able to mono- and dimethylate histone H1-4 at 'Lys-26' (H1.4K26me1 and H1.4K26me2, respectively) (PubMed:[19144645](#)). In addition to the histone methyltransferase activity, also methylates non-histone proteins: mediates dimethylation of 'Lys-373' of p53/TP53 (PubMed:[20118233](#)). Also methylates CDYL, WIZ, ACIN1, DNMT1, HDAC1, ERCC6, KLF12 and itself (PubMed:[18438403](#)).

Cellular Location

Nucleus. Chromosome. Note=Associates with euchromatic regions (PubMed:11316813). Does not associate with heterochromatin (PubMed:11316813).

Tissue Location

Expressed in all tissues examined, with high levels in fetal liver, thymus, lymph node, spleen and peripheral blood leukocytes and lower level in bone marrow

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.