

KD-Validated Anti-ALKBH1 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI2365

Product Information

Application	WB, ICC
Primary Accession	Q13686
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	23GB1385
Calculated MW	43832
Gene Name	ALKBH1
Aliases	ALKBH1; AlkB Homolog 1, Histone H2A Dioxygenase; ABH; ALKBH; HABH; Alpha-Ketoglutarate-Dependent Dioxygenase ABH1; Alkylated DNA Repair Protein AlkB Homolog 1; DNA N6-Methyl Adenine Demethylase ALKBH1; MRNA N(3)-Methylcytidine Demethylase; DNA Oxidative Demethylase ALKBH1; Nucleic Acid Dioxygenase ALKBH1; DNA 6mA Demethylase; DNA Lyase ABH1; AlkB; ABH1; AlkB, Alkylation Repair Homolog 1 (E. Coli); AlkB, Alkylation Repair Homolog (E. Coli); TRNA N1-Methyl Adenine Demethylase; AlkB, Alkylation Repair Homolog 1; Alkylation Repair, AlkB Homolog; DNA Demethylase ALKBH1; EC 1.14.11.51; EC 1.14.11.33; EC 1.14.11.-; EC 4.2.99.18; ALKB
Immunogen	A synthesized peptide derived from ALKBH1

Additional Information

Gene ID	8846
Other Names	Nucleic acid dioxygenase ALKBH1, 1.14.11.-, Alkylated DNA repair protein alkB homolog 1, Alpha-ketoglutarate-dependent dioxygenase ABH1, DNA 6mA demethylase, 1.14.11.-, ALKBH1 (HGNC:17911)

Protein Information

Name	ALKBH1 (HGNC:17911)
Function	Dioxygenase that acts on nucleic acids, such as DNA and tRNA (PubMed: 18603530 , PubMed: 27497299 , PubMed: 27745969). Requires molecular oxygen, alpha-ketoglutarate and iron (PubMed: 18603530 , PubMed: 27497299). A number of activities have been described for this dioxygenase, but recent results suggest that it mainly acts on tRNAs and mediates their demethylation or oxidation depending on the context and subcellular compartment (PubMed: 27497299 , PubMed: 27745969). Mainly acts as a tRNA demethylase by removing N(1)-methyladenine from various tRNAs, with a preference for N(1)-methyladenine at position 58 (m1A58) present on a stem loop structure of tRNAs (PubMed: 27745969). Acts as a regulator of

translation initiation and elongation in response to glucose deprivation: regulates both translation initiation, by mediating demethylation of tRNA(Met), and translation elongation, N(1)- methyladenine-containing tRNAs being preferentially recruited to polysomes to promote translation elongation (PubMed:[27745969](#)). In mitochondrion, specifically interacts with mt-tRNA(Met) and mediates oxidation of mt-tRNA(Met) methylated at cytosine(34) to form 5- formylcytosine (f(5)c) at this position (PubMed:[27497299](#)). mt-tRNA(Met) containing the f(5)c modification at the wobble position enables recognition of the AUA codon in addition to the AUG codon, expanding codon recognition in mitochondrial translation (PubMed:[27497299](#)). Specifically demethylates DNA methylated on the 6th position of adenine (N(6)-methyladenosine) DNA (PubMed:[30017583](#), PubMed:[30392959](#)). N(6)- methyladenosine (m6A) DNA is present at some L1 elements in embryonic stem cells and probably promotes their silencing (By similarity). Demethylates mRNAs containing N(3)-methylcytidine modification (PubMed:[31188562](#)). Also able to repair alkylated single-stranded DNA by oxidative demethylation, but with low activity (PubMed:[18603530](#)). Also has DNA lyase activity and introduces double-stranded breaks at abasic sites: cleaves both single-stranded DNA and double-stranded DNA at abasic sites, with the greatest activity towards double-stranded DNA with two abasic sites (PubMed:[19959401](#)). DNA lyase activity does not require alpha-ketoglutarate and iron and leads to the formation of an irreversible covalent protein-DNA adduct with the 5' DNA product (PubMed:[19959401](#), PubMed:[23577621](#)). DNA lyase activity is not required during base excision repair and class switch recombination of the immunoglobulin heavy chain during B lymphocyte activation. May play a role in placental trophoblast lineage differentiation (By similarity).

Cellular Location

Nucleus. Mitochondrion. Note=Mainly localizes in euchromatin, largely excluded from heterochromatin and nucleoli (By similarity).
{ECO:0000250|UniProtKB:P0CB42}

Tissue Location

Ubiquitous.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.