

KO-Validated Anti-SRPK2 Mouse Monoclonal Antibody

Mouse monoclonal antibody

Catalog # AGI2420

Product Information

Application	WB, FC
Primary Accession	P78362
Reactivity	Rat, Human
Clonality	Monoclonal
Isotype	Mouse IgG2a
Clone Names	25GB660
Calculated MW	77527
Gene Name	SRPK2
Aliases	SRPK2; SRSF Protein Kinase 2; SFRS Protein Kinase 2; SFRSK2; Serine/Arginine-Rich Protein-Specific Kinase 2; Serine/Arginine-Rich Splicing Factor Kinase 2; SR-Protein-Specific Kinase 2; SR Protein Kinase 2; EC 2.7.11.1; Serine/Threonine-Protein Kinase SRPK2; Serine Kinase SRPK2
Immunogen	Recombinant protein of human SRPK2

Additional Information

Gene ID	6733
Other Names	SRSF protein kinase 2, 2.7.11.1, SFRS protein kinase 2, Serine/arginine-rich protein-specific kinase 2, SR-protein-specific kinase 2, SRSF protein kinase 2 N-terminal, SRSF protein kinase 2 C-terminal, SRPK2 {ECO:0000312 EMBL:AAH68547.1}

Protein Information

Name	SRPK2 {ECO:0000312 EMBL:AAH68547.1}
Function	Serine/arginine-rich protein-specific kinase which specifically phosphorylates its substrates at serine residues located in regions rich in arginine/serine dipeptides, known as RS domains and is involved in the phosphorylation of SR splicing factors and the regulation of splicing (PubMed: 18559500 , PubMed: 21056976 , PubMed: 9472028). Promotes neuronal apoptosis by up-regulating cyclin-D1 (CCND1) expression (PubMed: 19592491). This is done by the phosphorylation of SRSF2, leading to the suppression of p53/TP53 phosphorylation thereby relieving the repressive effect of p53/TP53 on cyclin-D1 (CCND1) expression (PubMed: 21205200). Phosphorylates ACIN1, and redistributes it from the nuclear speckles to the nucleoplasm, resulting in cyclin A1 but not cyclin A2 up-regulation (PubMed: 18559500). Plays an essential role in spliceosomal B complex formation via the phosphorylation of DDX23/PRP28 (PubMed: 18425142). Probably by phosphorylating DDX23, leads to the suppression of incorrect R-loops formed during transcription; R-loops are composed of a DNA:RNA hybrid and the associated non-template

single-stranded DNA (PubMed:[28076779](#)). Can mediate hepatitis B virus (HBV) core protein phosphorylation (PubMed:[12134018](#)). Plays a negative role in the regulation of HBV replication through a mechanism not involving the phosphorylation of the core protein but by reducing the packaging efficiency of the pregenomic RNA (pgRNA) without affecting the formation of the viral core particles (PubMed:[16122776](#)). Phosphorylates the N-terminus of ERC1 (By similarity).

Cellular Location

Cytoplasm. Nucleus, nucleoplasm. Nucleus speckle. Chromosome.
Note=Shuttles between the nucleus and the cytoplasm (PubMed:19592491, PubMed:21056976, PubMed:21157427) KAT5/TIP60 inhibits its nuclear translocation (PubMed:21157427) Phosphorylation at Thr-492 by PKB/AKT1 promotes nuclear translocation (PubMed:19592491). Preferentially localizes across the entire gene coding region (PubMed:28076779). During transcription, accumulates at chromatin loci where unscheduled R-loops form and colocalizes with paused 'Ser-5'-phosphorylated POLR2A/RNA polymerase II and helicase DDX23 (PubMed:28076779).

Tissue Location

Highly expressed in brain, moderately expressed in heart and skeletal muscle and at low levels in lung, liver, and kidney

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.