

# CD45RB Antibody [Clone BRA-11; same as BRA-11G]

Purified Mouse Monoclonal Antibody

Catalog # AH10088

## Product Information

Application	FC
Primary Accession	<a href="#">P08575</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Names	BRA-11; same as BRA-11G
Calculated MW	147486

## Additional Information

Gene ID	5788
Other Names	Receptor-type tyrosine-protein phosphatase C, Leukocyte common antigen, L-CA, T200, CD45, PTPRC, CD45
Target/Specificity	Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)
Application Note	Flow Cytometry 2.5ul (0.5ug) per test per one million cells.
Format	0.5 ml at 200ug/ml; Conjugated to PE
Storage	Store at 2 to 8°C. Antibody is stable for 24 months.
Precautions	CD45RB Antibody [Clone BRA-11; same as BRA-11G] is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Name	PTPRC ( <a href="#">HGNC:9666</a> )
Synonyms	CD45
Function	Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor (PubMed: <a href="#">35767951</a> ). Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN. Dephosphorylates LYN, and thereby modulates LYN activity (By similarity). Interacts with CLEC10A at antigen presenting cell-T cell contact; CLEC10A on immature dendritic cells recognizes Tn antigen- carrying

PTPRC/CD45 receptor on effector T cells and modulates T cell activation threshold to limit autoreactivity.

**Cellular Location**

Cell membrane; Single-pass type I membrane protein. Membrane raft. Synapse. Note=Colocalized with DPP4 in membrane rafts.

**Tissue Location**

Isoform 1: Detected in thymocytes. Isoform 2: Detected in thymocytes. Isoform 3: Detected in thymocytes. Isoform 4: Not detected in thymocytes. Isoform 5: Detected in thymocytes. Isoform 6: Not detected in thymocytes. Isoform 7: Detected in thymocytes. Isoform 8: Not detected in thymocytes.

## Background

---

CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.


## References

---

1. Sedlak et al. Neoplasma 1989; 643, 1989.
2. Sutherland et.al Int. Immunol., 14(8): 953 – 962 (August 1, 2002)
3. Lim et.al Int. Immunol., 18(2):291-300 (2006)

## Images

---



Formalin-fixed, paraffin-embedded human tonsil stained with CD45RB Ab (BRA-11).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.