

CD35 / CR1 (Follicular Dendritic Cell Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SPM554]
Catalog # AH10415

Product Information

Application	IF, FC, IHC-P
Primary Accession	P17927
Other Accession	1378 , 334019
Reactivity	Human, Rhesus, Baboon, Cynomolgus
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Clone Names	SPM554
Calculated MW	223663

Additional Information

Gene ID	1378
Other Names	Complement receptor type 1, C3b/C4b receptor, CD35, CR1, C3BR
Application Note	IF~~1:50~200 FC~~1:10~50 IHC-P~~N/A
Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage	Store at 2 to 8°C. Antibody is stable for 24 months.
Precautions	CD35 / CR1 (Follicular Dendritic Cell Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CR1
Synonyms	C3BR
Function	Membrane immune adherence receptor that plays a critical role in the capture and clearance of complement-opsonized pathogens by erythrocytes and monocytes/macrophages (PubMed: 2963069). Mediates the binding by these cells of particles and immune complexes that have activated complement to eliminate them from the circulation (PubMed: 2963069). Also acts in the inhibition of spontaneous complement activation by impairing the

formation and function of the alternative and classical pathway C3/C5 convertases, and by serving as a cofactor for the cleavage by factor I of C3b to iC3b, C3c and C3d,g, and of C4b to C4c and C4d (PubMed:[2972794](#), PubMed:[8175757](#)). Also plays a role in immune regulation by contributing, upon ligand binding, to the generation of regulatory T cells from activated helper T cells (PubMed:[25742728](#)).

Cellular Location

Membrane; Single-pass type I membrane protein

Tissue Location

Present on erythrocytes, a subset of T cells, mature B cells, follicular dendritic cells, monocytes and granulocytes


Background

Recognizes a protein of 210-220kDa, which is identified as the complement receptor 1 (CR1)/CD35. This MAb is specific for a site in CR1 that is distal from the C3b-binding site, so that it is unable to block CR1 activity. This MAb is highly specific to CR1 and shows no cross-reaction with CR2. The primary function of CR1 is to serve as the cellular receptor for C3b and C4b, the most important components of the complement system leading to clearance of foreign macromolecules. The Knops blood group system is a system of antigens located on this protein. Follicular dendritic cells (FDC) are restricted to the B-cell regions of secondary lymphoid follicles. They are CD21+/CD35+/CD1a-. This MAb labels follicular dendritic cells and follicular dendritic cell sarcoma.

References

Hogg N, et. al. European Journal of Immunology, 1984, 14:236-43

Images



Formalin-fixed, paraffin-embedded human Tonsil stained with CD35 Monoclonal Antibody (SPM554).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.