

IgG (Immunoglobulin Gamma Heavy Chain) (B-Cell Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone IG217 + IG266]
Catalog # AH10501

Product Information

Application	IF, FC, IHC-P
Primary Accession	P01857
Other Accession	3500 (IGHG1) , 3501 (IGHG2) , 3502 (IGHG3) , 3503 (IGHG4) , 510635 , P01859 , P01860 , P01861
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1 + IgG2a, kappa
Clone Names	IG217 + IG266
Calculated MW	43912

Additional Information

Other Names	Ig gamma-1 chain C region, IGHG1
Application Note	IF~~1:50~200 FC~~1:10~50 IHC-P~~N/A
Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage	Store at 2 to 8°C. Antibody is stable for 24 months.
Precautions	IgG (Immunoglobulin Gamma Heavy Chain) (B-Cell Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	IGHG1 {ECO:0000303 PubMed:11340299, ECO:0000303 Ref.14}
Function	Constant region of immunoglobulin (Ig) heavy chains. Igs are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound Igs serve as receptors, which upon binding to a specific antigen trigger the clonal expansion and differentiation of B lymphocytes into Ig-secreting plasma cells. Secreted Igs known as antibodies mediate the effector phase of humoral immunity by blocking the interaction of infectious antigens with cellular receptors (via the antigen-binding region) and eliciting effector mechanisms that lead to pathogen neutralization (via the constant region)


(PubMed:[17576170](#), PubMed:[20176268](#), PubMed:[22158414](#)). The antigen-binding region is formed by the variable domain of one heavy chain paired with the variable domain of its associated light chain. Each Ig molecule has two antigen-binding sites with remarkable affinity for a particular antigen due to V-(D)-J rearrangement, somatic hypermutations and affinity maturation of the variable domains upon antigen exposure (PubMed:[17576170](#), PubMed:[20176268](#), PubMed:[22158414](#)). The constant region defines the Ig isotype that perform distinct sets of effector functions. B cells diversify and rearrange their Ig constant regions through class-switch recombination, a process by which the constant region is switched from one Ig isotype to another, namely from IgM and IgD to IgG, IgA and IgE (PubMed:[17576170](#), PubMed:[20176268](#), PubMed:[22158414](#)). The constant region of Ig gamma-1 (IgG1) isotype interacts (via the fragment crystallizable, Fc) with receptors on innate immune cells and the complement system to mediate humoral effector functions, including antibody-dependent cellular cytotoxicity or phagocytosis, complement- dependent cytotoxicity and inflammatory responses.

Cellular Location [Isoform 1]: Secreted

Background

This MAb is specific to heavy chain of IgG and shows minimal cross-reaction with heavy chains of other immunoglobulins. It is reactive with all subclasses of γ heavy chain (IgG1, IgG2a, IgG2b, IgG3). The most common feature of plasmacytomas, and certain non-Hodgkin Σ lymphomas is the restricted expression of a single heavy chain class. Demonstration of clonality in lymphoid infiltrates indicates that the infiltrate is clonal and therefore malignant. Ab-3 is superb for staining of formalin-fixed tissues.

Images



Formalin-fixed, paraffin-embedded human Tonsil stained with IgG Monoclonal Antibody (IG217 + IG266)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.