

# Insulin / IRDN (beta-Cell & Insulinoma Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SPM139 ]  
Catalog # AH10530

## Product Information

---

<b>Application</b>	IF, FC, IHC-P
<b>Primary Accession</b>	<a href="#">P01308</a>
<b>Other Accession</b>	<a href="#">3630</a> , <a href="#">272259</a>
<b>Reactivity</b>	Human, Pig, Bovine
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	Mouse / IgG1, kappa
<b>Clone Names</b>	SPM139
<b>Calculated MW</b>	11981

## Additional Information

---

<b>Gene ID</b>	3630
<b>Other Names</b>	Insulin, Insulin B chain, Insulin A chain, INS
<b>Application Note</b>	IF~~1:50~200 FC~~1:10~50 IHC-P~~N/A
<b>Format</b>	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage</b>	Store at 2 to 8°C.Antibody is stable for 24 months.
<b>Precautions</b>	Insulin / IRDN (beta-Cell & Insulinoma Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	INS
<b>Function</b>	Insulin decreases blood glucose concentration. It increases cell permeability to monosaccharides, amino acids and fatty acids. It accelerates glycolysis, the pentose phosphate cycle, and glycogen synthesis in liver.
<b>Cellular Location</b>	Secreted.

## Background

---

Recognizes a polypeptide which is identified as insulin, a 51-amino acid polypeptide composed of A and B chains connected through the C-peptide. Proinsulin, which has very little biological activity, is cleaved by proteases within its cell of origin into the insulin molecule and the C-terminal basic residue. Insulin enhances membrane transport of glucose, amino acids, and certain ions. It also promotes glycogen storage, formation of triglycerides, and synthesis of proteins and nucleic acids. Deficiency of insulin results in diabetes mellitus. The main storage site for insulin is the pancreatic islets. Antibodies to insulin are important as beta-cell and insulinoma marker.


## References

---

de la Tour, D., et al. 2001. Mol. Endoc. 15: 476-483. | Rajagopal, J., et al. 2003. Science 299: 363. | Morisset, J., et al. 2003. J. Histochem. Cytochem. 51: 1501-1513

## Images

---



Formalin-fixed, paraffin-embedded human Pancreas stained with Insulin Monoclonal Antibody (SPM139).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.