

# NKX2.2 (Neuroendocrine & Ewing Sarcoma Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SPM564] Catalog # AH10640

## **Product Information**

Application IF, FC, IHC-P
Primary Accession 095096
Other Accession 4821, 516922

Reactivity Human, Mouse, Chicken

**Host** Mouse **Clonality** Monoclonal

**Isotype** Mouse / IgG2b, kappa

Clone Names SPM564 Calculated MW 30133

#### Additional Information

**Gene ID** 4821

Other Names Homeobox protein Nkx-2.2, Homeobox protein NK-2 homolog B, NKX2-2,

NKX2.2, NKX2B

**Application Note** IF~~1:50~200 FC~~1:10~50 IHC-P~~N/A

**Format** 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G.

Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available

WITHOUT BSA & azide at 1.0mg/ml.

**Storage** Store at 2 to 8°C.Antibody is stable for 24 months.

**Precautions** NKX2.2 (Neuroendocrine & Ewing Sarcoma Marker) Antibody - With BSA and

Azide is for research use only and not for use in diagnostic or therapeutic

procedures.

#### **Protein Information**

Name NKX2-2

Synonyms NKX2.2, NKX2B

**Function** Transcriptional activator involved in the development of insulin-producting

beta cells in the endocrine pancreas (By similarity). May also be involved in specifying diencephalic neuromeric boundaries, and in controlling the expression of genes that play a role in axonal guidance. Binds to elements

within the NEUROD1 promoter (By similarity).


# **Background**

Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein specific to Ewing sarcoma, and was shown to be differentially upregulated in Ewing sarcoma on the basis of array-based gene expression analysis. It acts as a valuable marker for Ewing sarcoma, with a sensitivity of 93% and a specificity of 89%, and aids in the differential diagnosis of small round cell tumors.


### References

Wang, Y.C. et al. (2009) Endocr Relat Cancer 16, 267-79. | Yoshida A et. al. Am J Surg Pathol. 2012;36(7):993-9

## **Images**



Formalin-fixed paraffin-embedded human Ewing Is sarcoma stained with NKX2.2 Monoclonal Antibody (SPM564).



Formalin-fixed paraffin-embedded human Pancreas stained with NKX2.2 Monoclonal Antibody (SPM564).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.