

# CD45RA (Leucocyte Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SPM568 ]  
Catalog # AH10691

## Product Information

---

|                   |   |
|-------------------|---|
| Application       | IF, FC, IHC-P                                 |
| Primary Accession | <a href="#">P08575</a>                        |
| Other Accession   | <a href="#">5788</a> , <a href="#">654514</a> |
| Reactivity        | Human   |
| Host              | Mouse   |
| Clonality         | Monoclonal                                    |
| Isotype           | Mouse / IgG2a, kappa                          |
| Clone Names       | SPM568  |
| Calculated MW     | 147486  |

## Additional Information

---

|                  |  |
|------------------|--|
| Gene ID          | 5788   |
| Other Names      | Receptor-type tyrosine-protein phosphatase C, 3.1.3.48, Leukocyte common antigen, L-CA, T200, CD45, PTPRC, CD45  |
| Application Note | IF~~1:50~200 FC~~1:10~50 IHC-P~~N/A  |
| Format           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Storage          | Store at 2 to 8°C. Antibody is stable for 24 months.   |
| Precautions      | CD45RA (Leucocyte Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.                              |

## Protein Information

---

|          |  |
|----------|--|
| Name     | PTPRC ( <a href="#">HGNC:9666</a> )  |
| Synonyms | CD45   |
| Function | Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor (PubMed: <a href="#">35767951</a> ). Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN. Dephosphorylates LYN, and thereby modulates LYN activity (By |

similarity). Interacts with CLEC10A at antigen presenting cell-T cell contact; CLEC10A on immature dendritic cells recognizes Tn antigen- carrying PTPRC/CD45 receptor on effector T cells and modulates T cell activation threshold to limit autoreactivity.

**Cellular Location**

Cell membrane; Single-pass type I membrane protein. Membrane raft. Synapse. Note=Colocalized with DPP4 in membrane rafts.

**Tissue Location**

Isoform 1: Detected in thymocytes. Isoform 2: Detected in thymocytes. Isoform 3: Detected in thymocytes. Isoform 4: Not detected in thymocytes. Isoform 5: Detected in thymocytes. Isoform 6: Not detected in thymocytes. Isoform 7: Detected in thymocytes Isoform 8: Not detected in thymocytes.

## Background

---

Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C, respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% of peripheral CD8+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory T-cells. CD45RA and CD45RO define complementary, predominantly non-overlapping populations of resting peripheral T-cells. This MAb is useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especially be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphomas (CD45RA +ve).


## References

---

Oxford University Press, Oxford, p511-515, 1995. | Clement LT. J Clin Immunol. 1992, 12(1):1-10. | Yamada A, et al. Cell Immunol. 1992, 142(1):114-24. | Mahalingam M, et al. Clin Immunol Immunopathol. 1996,81(2):210-214. Jacob MC, et al. Am J Hematol. 1992, 39(1):45-5

## Images

---



Formalin-fixed, paraffin-embedded human Tonsil stained with CD45RA Monoclonal Antibody (SPM568).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.