

# Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SPM117 ]  
Catalog # AH10717

## Product Information

---

Application	WB, IF, FC, IHC-P
Primary Accession	<a href="#">P10415</a>
Other Accession	<a href="#">596</a> , <a href="#">150749</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Clone Names	SPM117
Calculated MW	26266

## Additional Information

---

Gene ID	596
Other Names	Apoptosis regulator Bcl-2, BCL2
Application Note	WB~~1:1000 IF~~1:50~200 FC~~1:10~50 IHC-P~~N/A
Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage	Store at 2 to 8°C. Antibody is stable for 24 months.
Precautions	Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	BCL2
Function	Suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural cells (PubMed: <a href="#">1508712</a> , PubMed: <a href="#">8183370</a> ). Regulates cell death by controlling the mitochondrial membrane permeability (PubMed: <a href="#">11368354</a> ). Appears to function in a feedback loop system with caspases (PubMed: <a href="#">11368354</a> ). Inhibits caspase activity either by preventing the release of cytochrome c from the mitochondria and/or by binding to the apoptosis-activating factor (APAF-1) (PubMed: <a href="#">11368354</a> ). Also acts as an inhibitor of autophagy: interacts with BECN1 and AMBRA1 during

non-starvation conditions and inhibits their autophagy function (PubMed:[18570871](#), PubMed:[20889974](#), PubMed:[21358617](#)). May attenuate inflammation by impairing NLRP1- inflammasome activation, hence CASP1 activation and IL1B release (PubMed:[17418785](#)).

#### Cellular Location

Mitochondrion outer membrane; Single-pass membrane protein. Nucleus membrane; Single-pass membrane protein. Endoplasmic reticulum membrane; Single-pass membrane protein. Cytoplasm {ECO:0000250|UniProtKB:P10417}

#### Tissue Location

Expressed in a variety of tissues.

## Background

---

This antibody recognizes a protein of 25-26kDa, identified as the bcl-2  $\alpha$  oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of bcl-2  $\alpha$  oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers express high levels of bcl-2  $\alpha$  protein, whereas the normal or hyperplastic germinal centers are negative. Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be used in distinguishing between those follicular lymphomas that express bcl-2 protein and the small number in which the neoplastic cells are bcl-2 negative.


## References

---


Pezzella F et. al.. American Journal of Pathology, 1990, 137(2):225-32

## Images

---




Formalin-fixed, paraffin-embedded human Non-Hodgkin's Lymphoma stained with Bcl-2 Ab (SPM117).



Flow Cytometry of Jurkat cells using Bcl-2 Ab (SPM117) (red) and isotype control (green).

Western Blot of Bcl-2 in human Skin using Bcl-2 Ab (SPM117).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.