

# von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SPM577 ]  
Catalog # AH10813

## Product Information

---

Application	WB, IF, FC, IP, IHC-P
Primary Accession	<a href="#">P04275</a>
Other Accession	<a href="#">7450</a> , <a href="#">440848</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Clone Names	SPM577
Calculated MW	309265

## Additional Information

---

Gene ID	7450
Other Names	von Willebrand factor, vWF, von Willebrand antigen 2, von Willebrand antigen II, VWF, F8VWF
Application Note	WB~~1:1000 IF~~1:50~200 FC~~1:10~50 IP~~N/A IHC-P~~N/A
Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage	Store at 2 to 8°C. Antibody is stable for 24 months.
Precautions	von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	VWF
Synonyms	F8VWF
Function	Important in the maintenance of hemostasis, it promotes adhesion of platelets to the sites of vascular injury by forming a molecular bridge between sub-endothelial collagen matrix and platelet- surface receptor complex GPIb-IX-V. Also acts as a chaperone for coagulation factor VIII, delivering it to the site of injury, stabilizing its heterodimeric structure and protecting it from

premature clearance from plasma.

**Cellular Location**

Secreted. Secreted, extracellular space, extracellular matrix. Note=Localized to storage granules

**Tissue Location**

Plasma.

## Background

---

von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi's sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.


## References

---

Motta, A. et al. 2009. J Biomater Sci Polym Ed. 20: 1875-1897. | Germann, B. et al. 2008. Pharmazie. 63: 303-307

## Images

---



Formalin-fixed, paraffin-embedded human Tonsil stained with vWF Monoclonal Antibody (SPM577)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.