

# Glycophorin A / CD235a (Erythrocyte Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone GYPA/280] Catalog # AH11379

#### **Product Information**

Application IHC, IF, FC Primary Accession P02724

Other Accession 2993, 2994, 434973, 654368

Reactivity Human
Host Mouse
Clonality Monoclonal

Isotype Mouse / IgG1, kappa

Clone Names GYPA/280 Calculated MW 16430

## **Additional Information**

**Gene ID** 2993

Other Names Glycophorin-A, MN sialoglycoprotein, PAS-2, Sialoglycoprotein alpha, CD235a,

GYPA, GPA

**Application Note** IHC~~1:100~500 IF~~1:50~200 FC~~1:10~50

**Storage** Store at 2 to 8°C.Antibody is stable for 24 months.

**Precautions** Glycophorin A / CD235a (Erythrocyte Marker) Antibody - With BSA and Azide

is for research use only and not for use in diagnostic or therapeutic

procedures.

#### **Protein Information**

Name GYPA ( HGNC:4702)

**Function** Component of the ankyrin-1 complex, a multiprotein complex involved in

the stability and shape of the erythrocyte membrane (PubMed: 35835865). Glycophorin A is the major intrinsic membrane protein of the erythrocyte. The

N-terminal glycosylated segment, which lies outside the erythrocyte

membrane, has MN blood group receptors. Appears to be important for the function of SLC4A1 and is required for high activity of SLC4A1. May be

involved in translocation of SLC4A1 to the plasma membrane.

Cellular Location Cell membrane; Single-pass type I membrane protein Note=Appears to be

colocalized with SLC4A1


# **Background**

Recognizes a sialoglycoprotein of 39kDa, identified as glycophorin A (GPA). It is present on red blood cells (RBC) and erythroid precursor cells. It has been shown that glycophorin acts as the receptor for Sandei virus and parvovirus. Glycophorins A (GPA) and B (GPB), which are single, trans-membrane sialoglycoproteins. GPA is the carrier of blood group M and N specificities, while GPB accounts for S and U specificities. GPA and GPB provide the cells with a large mucin like surface and it has been suggested this provides a barrier to cell fusion, so minimizing aggregation between red blood cells in the circulation.


### References

Andersson, L.C., et al. 1979. Glycophorin A as a cell surface marker of early erythroid differentiation in acute leukemia. Int. J. Cancer 23: 717-720. | Liszka, K., et al., 1983. Glycophorin A expression in malignant hematopoiesis. Am. J. Hematol. 15: 219-226

# **Images**



Formalin-fixed, paraffin-embedded human Placenta stained with Glycophorin A Monoclonal Antibody (GYPA/280)



Formalin-fixed, paraffin-embedded human Angiosarcoma stained with Glycophorin A Monoclonal Antibody (GYPA/280)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.