

# CD50 / ICAM-3 Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone ICAM3/1019] Catalog # AH11492

# **Product Information**

Application Primary Accession	WB, IHC, IF, FC P32942
Other Accession	<u>r 32942</u> <u>3385, 354563</u>
Reactivity	<u>5585</u> , <u>554505</u> Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG2b, kappa
Clone Names	ICAM3/1019
Calculated MW	59541

## **Additional Information**

Gene ID	3385
Other Names	Intercellular adhesion molecule 3, ICAM-3, CDw50, ICAM-R, CD50, ICAM3
Application Note	WB~~1:1000 IHC~~1:100~500 IF~~1:50~200 FC~~1:10~50
Storage	Store at 2 to 8°C.Antibody is stable for 24 months.
Precautions	CD50 / ICAM-3 Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

#### **Protein Information**

Name	ICAM3
Function	ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2) (PubMed: <u>1448173</u> ). ICAM3 is also a ligand for integrin alpha-D/beta-2. In association with integrin alpha- L/beta-2, contributes to apoptotic neutrophil phagocytosis by macrophages (PubMed: <u>23775590</u> ).
Cellular Location	Membrane; Single-pass type I membrane protein.
Tissue Location	Leukocytes.

## Background

Recognizes an N-glycosylated glycoprotein of 120kDa with intra-chain disulfide bonds, identified as CD50 or ICAM-3. CD50 is the major ligand for LFA-1 (CD11a/CD18) and may have signaling role to increase adhesion.

It is expressed on thymocytes and T lymphocytes and is resistant to treatment with phosphatidylinositol (PI) phospholipase C. This MAb is excellent for staining of formalin/paraffin tissues.

## References

Fawcett, J., Holness, C.L., Needham, L.A., Turley, H., Gatter, K.C., Mason, D.Y. and Simmons, D.L. 1992. Molecular cloning of ICAM-3, a third ligand for LFA-1, constitutively expressed on resting leukocytes. Nature 360:481-484

#### Images



Western Blot Analysis of HL-60 Cell Lysate using CD50 Monoclonal Antibody (ICAM3/1019)



Formalin-fixed, paraffin-embedded human Tonsil stained with CD50 Monoclonal Antibody (ICAM3/1019)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.