

# Cytokeratin 17 (KRT17) (Basal Epithelial Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone KRT17/778 ]

Catalog # AH11695

## Product Information

---

<b>Application</b>	IHC, IF, FC
<b>Primary Accession</b>	<a href="#">Q04695</a>
<b>Other Accession</b>	<a href="#">3872</a> , <a href="#">2785</a>
<b>Reactivity</b>	Human, Rat, Pig, Goat, Bovine
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	Mouse / IgG2b, kappa
<b>Clone Names</b>	KRT17/778
<b>Calculated MW</b>	48106

## Additional Information

---

<b>Gene ID</b>	3872
<b>Other Names</b>	Keratin, type I cytoskeletal 17, 39.1, Cytokeratin-17, CK-17, Keratin-17, K17, KRT17
<b>Application Note</b>	IHC~~1:100~500 IF~~1:50~200 FC~~1:10~50
<b>Storage</b>	Store at 2 to 8°C.Antibody is stable for 24 months.
<b>Precautions</b>	Cytokeratin 17 (KRT17) (Basal Epithelial Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	KRT17
<b>Function</b>	Type I keratin involved in the formation and maintenance of various skin appendages, specifically in determining shape and orientation of hair (By similarity). Required for the correct growth of hair follicles, in particular for the persistence of the anagen (growth) state (By similarity). Modulates the function of TNF-alpha in the specific context of hair cycling. Regulates protein synthesis and epithelial cell growth through binding to the adapter protein SFN and by stimulating Akt/mTOR pathway (By similarity). Involved in tissue repair. May be a marker of basal cell differentiation in complex epithelia and therefore indicative of a certain type of epithelial 'stem cells'. Acts as a promoter of epithelial proliferation by acting a regulator of immune response in skin: promotes Th1/Th17-dominated immune environment contributing to

the development of basaloid skin tumors (By similarity). May act as an autoantigen in the immunopathogenesis of psoriasis, with certain peptide regions being a major target for autoreactive T-cells and hence causing their proliferation.

**Cellular Location**

Cytoplasm {ECO:0000250|UniProtKB:Q9QWL7}.

**Tissue Location**

Expressed in the outer root sheath and medulla region of hair follicle specifically from eyebrow and beard, digital pulp, nail matrix and nail bed epithelium, mucosal stratified squamous epithelia and in basal cells of oral epithelium, palmoplantar epidermis and sweat and mammary glands. Also expressed in myoepithelium of prostate, basal layer of urinary bladder, cambial cells of sebaceous gland and in exocervix (at protein level)

## Background

---

Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excellent tool to distinguish myoepithelial cells from luminal epithelium of various glands such as mammary, sweat and salivary. CK17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. It may be considered as an epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in SCLC much higher than in LADC. Eighty-five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course. The histologic differentiation of ampullary cancer, intestinal vs. pancreatobiliary, is very important for treatment. Usually anti-CK17 and anti-MUC1 immunoreactivity represents pancreatobiliary subtype whereas anti-MUC2 and anti-CDX-2 positivity defines intestinal subtype.


## References

---


(1) Smedts et. al. Am J Pathol 141: 497, 1992. (2) Smedts et. al. Am J Pathol 140: 601, 1992.(3) Wetzels et. al. Histopathol 20: 295, 1992. |

## Images

---



Formalin-fixed, paraffin-embedded human Bladder Carcinoma stained with CK17 Monoclonal Antibody (KRT17/778).



Formalin-fixed, paraffin-embedded human Cervical Carcinoma stained with CK17 Monoclonal Antibody (KRT17/778).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.