

CD99 / MIC2 (Ewing's Sarcoma Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SPM586]

Catalog # AH11818

Product Information

Application	IHC, IF, FC
Primary Accession	P14209
Other Accession	4267 , 653349
Reactivity	Human, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgM, kappa
Clone Names	SPM586
Calculated MW	18848

Additional Information

Gene ID	4267
Other Names	CD99 antigen, 12E7, E2 antigen, Protein MIC2, T-cell surface glycoprotein E2, CD99, CD99, MIC2, MIC2X, MIC2Y
Application Note	IHC~~1:100~500 IF~~1:50~200 FC~~1:10~50
Storage	Store at 2 to 8°C.Antibody is stable for 24 months.
Precautions	CD99 / MIC2 (Ewing's Sarcoma Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CD99
Synonyms	MIC2, MIC2X, MIC2Y
Function	Involved in T-cell adhesion processes and in spontaneous rosette formation with erythrocytes. Plays a role in a late step of leukocyte extravasation helping leukocytes to overcome the endothelial basement membrane. Acts at the same site as, but independently of, PECAM1. Involved in T-cell adhesion processes (By similarity).
Cellular Location	Membrane; Single-pass type I membrane protein


Background

Recognizes a sialoglycoprotein of 27-32kDa, identified as CD99, or MIC2 gene product, or E2 antigen. This antigen is expressed on the cell membrane of some lymphocytes, cortical thymocytes, and granulosa cells of the ovary. Most pancreatic islet cells, Sertoli cells of the testis, and some endothelial cells express this antigen. Mature granulocytes express very little or no CD99. MIC2 is strongly expressed on Ewing's sarcoma cells and primitive peripheral neuroectodermal tumors. This MAb shows a very similar reactivity to other CD99 MAbs (e.g. O13, 12E7, or HBA-71) and is excellent for immunohistochemical staining of formalin-fixed, paraffin-embedded tissues.

References

Sandrin MS, et. al. Immunogenetics, 1992, 35(4):283-5

Images



Formalin-fixed, paraffin-embedded human Ewing's sarcoma stained with CD99 Monoclonal Antibody (SPM586).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.