

# CD99 / MIC2 (Ewings Sarcoma Marker) Antibody - Culture Supernatant

Mouse Monoclonal Antibody [Clone 12E7.]  
Catalog # AH11820

## Product Information

---

|                          |   |
|--------------------------|---|
| <b>Application</b>       | IHC, IF, FC                                   |
| <b>Primary Accession</b> | <a href="#">P14209</a>                        |
| <b>Other Accession</b>   | <a href="#">4267</a> , <a href="#">653349</a> |
| <b>Reactivity</b>        | Human   |
| <b>Host</b>              | Mouse   |
| <b>Clonality</b>         | Monoclonal                                    |
| <b>Isotype</b>           | Mouse / IgG1, kappa                           |
| <b>Clone Names</b>       | 12E7.   |
| <b>Calculated MW</b>     | 27 KDa  |

## Additional Information

---

|                         |  |
|-------------------------|--|
| <b>Other Names</b>      | CD99 antigen, 12E7, E2 antigen, Protein MIC2, T-cell surface glycoprotein E2, CD99, CD99, MIC2, MIC2X, MIC2Y   |
| <b>Application Note</b> | IHC~~1:100~500 IF~~1:50~200 FC~~1:10~50  |
| <b>Storage</b>          | Store at 2 to 8°C. Antibody is stable for 24 months.   |
| <b>Precautions</b>      | CD99 / MIC2 (Ewings Sarcoma Marker) Antibody - Culture Supernatant is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

---

### Background

---

Recognizes a sialoglycoprotein of 27-32kDa, identified as CD99, or MIC2 gene product, or E2 antigen. MIC2 gene is located in the pseudo-autosomal region of the human X and Y chromosome. MIC2 gene encodes two distinct proteins, which are produced by alternative splicing of the CD99 gene transcript and are identified as bands of 30 and 32kDa (p30/32). Although its function is not fully understood, CD99 is implicated in various cellular processes including homotypic aggregation of T cells, upregulation of T cell receptor and MHS molecules, apoptosis of immature thymocytes and leukocyte diapedesis. CD99 is expressed on the cell membrane of some lymphocytes, cortical thymocytes, and granulosa cells of the ovary. Most pancreatic islet cells, Sertoli cells of the testis, and some endothelial cells express this antigen. Mature granulocytes express very little or no CD99. MIC2 is strongly expressed on Ewing s sarcoma cells and primitive peripheral neuroectodermal tumors.


## References

---

Levy R, Dilley J, Fox RI, Warnke R. A human thymus-leukemia antigen defined by hybridoma monoclonal antibodies. PNAS USA 1979;76(12):6552-6 |

## Images

---



Formalin-fixed, paraffin-embedded human Ewing's Sarcoma stained with CD99 Monoclonal Antibody (12E7).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.