

# MUC5AC (Mucin 5AC / Gastric Mucin) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone 58M1 ]  
Catalog # AH11904

## Product Information

---

|                   |   |
|-------------------|---|
| Application       | IHC, IF, FC                                   |
| Primary Accession | <a href="#">P98088</a>                        |
| Other Accession   | <a href="#">4586</a> , <a href="#">534332</a> |
| Reactivity        | Human   |
| Host              | Mouse   |
| Clonality         | Monoclonal                                    |
| Isotype           | Mouse / IgG1, kappa                           |
| Clone Names       | 58M1  |
| Calculated MW     | 585570  |

## Additional Information

---

|                  |   |
|------------------|---|
| Gene ID          | 4586  |
| Other Names      | Mucin-5AC, MUC-5AC, Gastric mucin, Lewis B blood group antigen, LeB, Major airway glycoprotein, Mucin-5 subtype AC, tracheobronchial, Tracheobronchial mucin, TBM, MUC5AC, MUC5 |
| Application Note | IHC~~1:100~500 IF~~1:50~200 FC~~1:10~50   |
| Storage          | Store at 2 to 8°C.Antibody is stable for 24 months.   |
| Precautions      | MUC5AC (Mucin 5AC / Gastric Mucin) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.                              |

## Protein Information

---

|                   |   |
|-------------------|---|
| Name              | MUC5AC {ECO:0000303   PubMed:11535137, ECO:0000312   HGNC:HGNC:7515}  |
| Function          | Gel-forming glycoprotein of gastric and respiratory tract epithelia that protects the mucosa from infection and chemical damage by binding to inhaled microorganisms and particles that are subsequently removed by the mucociliary system (PubMed: <a href="#">14535999</a> , PubMed: <a href="#">14718370</a> ). Interacts with H.pylori in the gastric epithelium, Barrett's esophagus as well as in gastric metaplasia of the duodenum (GMD) (PubMed: <a href="#">14535999</a> ). |
| Cellular Location | Secreted<br><br>Highly expressed in surface mucosal cells of respiratory tract and stomach  |

**Tissue Location**

epithelia. Overexpressed in a number of carcinomas. Also expressed in Barrett's esophagus epithelium and in the proximal duodenum.

**Background**

---

This MAb recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). □Its epitope is destroyed by beta-mercaptoethanol but not by periodate treatment. This mucin is present in primary ovarian mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies, Anti-MUC5AC may be useful for differential identification of primary mucinous ovarian tumors from colon adenocarcinoma metastatic to the ovary. MUC5AC antibodies may also be useful for identification of intestinal metaplasia as well as in the identification of pancreatic carcinoma and pre-cancerous changes vs. normal pancreas.


**References**

---

Cancer Res.46: 3983-3989 (1986). | Biochem. J. 254: 185-193 (1988). | Int. J. Cancer 47: 304-310 (1991). | J. Immunol. Methods 149: 105-113 (1992)

**Images**

---



Formalin-fixed, paraffin-embedded human Gastric Carcinoma stained with MUC5AC Monoclonal Antibody (58M1).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.