

# Nucleolin (Marker of Human Cells) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone 364-5 ]

Catalog # AH11984

## Product Information

---

Application	WB, IHC, IF, FC
Primary Accession	<a href="#">P19338</a>
Other Accession	<a href="#">4691</a> , <a href="#">79110</a>
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Clone Names	364-5
Calculated MW	76614

## Additional Information

---

Gene ID	4691
Other Names	Nucleolin, Protein C23, NCL
Application Note	WB~~1:1000 IHC~~1:100~500 IF~~1:50~200 FC~~1:10~50
Storage	Store at 2 to 8°C.Antibody is stable for 24 months.
Precautions	Nucleolin (Marker of Human Cells) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	NCL
Function	Nucleolin is the major nucleolar protein of growing eukaryotic cells. It is found associated with intranucleolar chromatin and pre-ribosomal particles. It induces chromatin decondensation by binding to histone H1. It is thought to play a role in pre-rRNA transcription and ribosome assembly. May play a role in the process of transcriptional elongation. Binds RNA oligonucleotides with 5'-UUAGGG- 3' repeats more tightly than the telomeric single-stranded DNA 5'- TTAGGG-3' repeats.
Cellular Location	Nucleus, nucleolus. Cytoplasm. Note=Localized in cytoplasmic mRNP granules containing untranslated mRNAs

## Background

---

Recognizes a protein of ~76kDa, which is identified as Nucleolin (NCL). It is the major nucleolar phosphoprotein of growing eukaryotic cells. NCL is located mainly in dense fibrillar regions of the nucleolus. It is found associated with intranucleolar chromatin and pre-ribosomal particles. Human NCL gene consists of 14 exons with 13 introns and spans approximately 11kb. It induces chromatin decondensation by binding to histone H1. It is thought to play a role in pre-rRNA transcription and ribosome assembly. □This MAb can be used to stain the nucleoli in cell or tissue preparations and can be used as a marker of the nucleoli in subcellular fractions. It produces a speckled pattern in the nuclei of cells of normal and malignant cells and may be used to stain the nucleoli of cells in fixed or frozen tissue sections. It can be used with paraformaldehyde fixed frozen tissue or cell preparations and formalin fixed, paraffin-embedded tissue sections.


## References

---


Fujiki H, Watanabe T, Suganuma M. Cell-surface nucleolin acts as a central mediator for carcinogenic, anti-carcinogenic, and disease-related ligands. *J Cancer Res Clin Oncol*. 2014;140(5):689-99. | Qiu W, Zhou F, Zhang Q, Sun X, Shi X, Liang Y, Wang X, Yue L. Overexpression of nucleolin and different expression sites both related to the prognosis of gastric cancer. *APMIS*. 2013;121(10):919-25.

## Images


---



Formalin-fixed paraffin-embedded human Skin stained with Nucleolin Monoclonal Antibody (364-5).




Formalin-fixed paraffin-embedded human Tonsil stained with Nucleolin Monoclonal Antibody (364-5).



Formalin-fixed paraffin-embedded human Colon Carcinoma stained with AF488 Conjugate of Nucleolin Monoclonal Antibody (364-5).

Flow Cytometry of Human Nucleolin Ag on 293T Cells.  
Black: Cells alone; Grey: Isotype Control; Green:  
AF488-labeled Nucleolin Monoclonal Antibody (364-5).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.