

# SOX10 (Melanoma Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SOX10/991] Catalog # AH12338

### **Product Information**

**Application** WB, IHC, IF, FC

Primary Accession
Other Accession
Reactivity
Host
Clonality
P56693
6663, 376984
Human
Mouse
Monoclonal

**Isotype** Mouse / IgG2b, kappa

Clone Names SOX10/991 Calculated MW 49911

#### Additional Information

Gene ID 6663

Other Names Transcription factor SOX-10, SOX10

**Application Note** WB~~1:1000 IHC~~1:100~500 IF~~1:50~200 FC~~1:10~50

**Storage** Store at 2 to 8°C.Antibody is stable for 24 months.

**Precautions** SOX10 (Melanoma Marker) Antibody - With BSA and Azide is for research use

only and not for use in diagnostic or therapeutic procedures.

#### **Protein Information**

Name SOX10

**Function** Transcription factor that plays a central role in developing and mature glia

(By similarity). Specifically activates expression of myelin genes, during oligodendrocyte (OL) maturation, such as DUSP15 and MYRF, thereby playing a central role in oligodendrocyte maturation and CNS myelination (By similarity). Once induced, MYRF cooperates with SOX10 to implement the myelination program (By similarity). Transcriptional activator of MITF, acting synergistically with PAX3 (PubMed:21965087). Transcriptional activator of

MBP, via binding to the gene promoter (By similarity).

**Cellular Location** Cytoplasm. Nucleus. Mitochondrion outer membrane

{ECO:0000250|UniProtKB:Q04888}; Peripheral membrane protein

{ECO:0000250 | UniProtKB:Q04888}; Cytoplasmic side

{ECO:0000250|UniProtKB:Q04888}


## **Background**

Recognizes a protein of ~55kDa, identified as SOX10. This MAb is highly specific and does not cross-react with other members of the SOX-family. SOX genes comprise a family of genes that are related to the mammalian sex-determining gene SRY. These genes similarly contain sequences that encode for the HMG-box domain, which is responsible for the sequence-specific DNA-binding activity. SOX-10 is a sensitive marker of melanoma, including conventional, spindled, and desmoplastic subtypes. It is expressed by metastatic melanomas and nodal capsular nevus in sentinel lymph nodes, but not by other lymph node components such as dendritic cells, which usually express S100 protein. Commonly used melanoma markers, such as anti-HMB-45 and anti-Melan-A, are poorly expressed in desmoplastic melanomas while SOX-10 is moderately to strongly expressed in desmoplastic melanomas. SOX-10 is considered as a very reliable marker for recognizing residual desmoplastic melanomas. In normal tissues, it is expressed in Schwann cells, melanocytes, and myoepithelial cells of salivary, bronchial and mammary glands. SOX-10 expression is also observed in mast cells.


#### References

Mohamed A, et al. SOX10 Expression in malignant melanoma, carcinoma, and normal tissues. Appl Immunohistochem Mol Morphol. 2013; 21(6):506-10.

## **Images**



Formalin-fixed, paraffin-embedded human Melanoma stained with SOX10 Monoclonal Antibody (SOX10/991).



Western Blot of SOX10 (A) Recombinant protein (B) A375 Cell Lysate using SOX10 Monoclonal Antibody (SOX10/991).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.