10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide

Rat Monoclonal Antibody [Clone HSP90B1/1192 ]
Catalog # AH12459

## Product Information

| | |
|---|---|
| **Application** | WB, IF, FC, IHC-P |
| **Primary Accession** | P14625 |
| **Other Accession** | 7184, 192374 |
| **Reactivity** | Human |
| **Host** | Rat |
| **Clonality** | Monoclonal |
| **Isotype** | Rat / IgG2a, kappa |
| **Clone Names** | HSP90B1/1192 |
| **Calculated MW** | 92469 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 7184 |
| **Other Names** | Endoplasmin, 94 kDa glucose-regulated protein, GRP-94, Heat shock protein 90 kDa beta member 1, Tumor rejection antigen 1, gp96 homolog, HSP90B1, GRP94, TRA1 |
| **Application Note** | WB~~1:1000 IF~~1:50~200 FC~~1:10~50 IHC-P~~N/A |
| **Storage** | Store at 2 to 8°C.Antibody is stable for 24 months. |
| **Precautions** | GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | HSP90B1 {ECO:0000303\|PubMed:39509507, ECO:0000312\|HGNC:HGNC:12028} |
| **Function** | ATP-dependent chaperone involved in the processing of proteins in the endoplasmic reticulum, regulating their transport (PubMed:23572575, PubMed:39509507). Together with MESD, acts as a modulator of the Wnt pathway by promoting the folding of LRP6, a coreceptor of the canonical Wnt pathway (PubMed:23572575, PubMed:39509507). When associated with CNPY3, required for proper folding of Toll-like receptors (PubMed:11584270). Promotes folding and trafficking of TLR4 to the cell surface (PubMed:11584270). May participate in the unfolding of cytosolic leaderless cargos (lacking the secretion signal sequence) such as the interleukin 1/IL-1 to |

| | |
|---|---|
| | facilitate their translocation into the ERGIC (endoplasmic reticulum- Golgi intermediate compartment) and secretion; the translocation process is mediated by the cargo receptor TMED10 (PubMed:32272059). |
| **Cellular Location** | Endoplasmic reticulum lumen. Sarcoplasmic reticulum lumen {ECO:0000250\|UniProtKB:P41148}. Melanosome Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV. |


## Background

Recognizes a protein of 94kDa, which is identified as the glucose-regulated protein 94 (grp94) and also tumor rejection antigen (gp96). Grp94 shows a high degree of sequence homology with the heat shock protein 90 (hsp90). This MAb is highly specific to grp94 and shows minimal cross-reaction with other members of the HSP90 family. Grp s are a class of proteins unresponsive to heat shock and are induced by glucose deprivation. Grp94 has been briefly studied as a prognostic factor in breast cancer.

## References

Sorger, P.K. et al. J. Mol. Biol. 194: 341-344 (1987). | Tandon, A.K. et.al. Breast Cancer Res. and Treat. 16: 146 (1990). |

## Images



Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with GRP94 Monoclonal Antibody (HSP90B1/1192).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.