

# CD70 / TNFSF7 (Activated T- & B-Lymphocyte Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone TNFS7/1026 ] Catalog # AH12802

# **Product Information**

Application Primary Accession	IF, FC, IP, E, IHC-F <u>P32970</u>
Other Accession	<u>970, 501497, 715224</u>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Clone Names	TNFS7/1026
Calculated MW	21118

## **Additional Information**

Gene ID	970
Other Names	CD70 antigen, CD27 ligand, CD27-L, Tumor necrosis factor ligand superfamily member 7, CD70, CD70, CD27L, CD27LG, TNFSF7
Application Note	IF~~1:50~200 FC~~1:10~50 IP~~N/A E~~N/A IHC-F~~N/A
Storage	Store at 2 to 8°C.Antibody is stable for 24 months.
Precautions	CD70 / TNFSF7 (Activated T- & B-Lymphocyte Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

#### **Protein Information**

Name	CD70 ( <u>HGNC:11937</u> )
Function	Expressed at the plasma membrane of B cells, it is the ligand of the CD27 receptor which is specifically expressed at the surface of T cells (PubMed: <u>28011863</u> , PubMed: <u>28011864</u> , PubMed: <u>8387892</u> ). The CD70- CD27 signaling pathway mediates antigen-specific T cell activation and expansion which in turn provides immune surveillance of B cells (PubMed: <u>28011863</u> , PubMed: <u>28011864</u> ).
Cellular Location	Cell membrane; Single-pass type II membrane protein

# Background

It recognizes a protein of 30kDa, identified as CD70. It is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T- and B-lymphocytes. It induces proliferation of co-stimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin synthesis.

## References

Lens, S.M., et al. 1995. CD27-CD70 interaction: unravelling its implication in normal and neoplastic B cell growth. Leuk. Lymphoma 18: 51-59

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.