

Cytokeratin, pan (Epithelial Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone KRTL/1077 + KRTH/1076]
Catalog # AH12903

Product Information

Application	IF, FC, IHC-P
Primary Accession	Q7Z794
Other Accession	374454 (KRT77) , 51350 (KRT76) , Q01546 (KRT76) Unigene334989 (KRT77) , 654392 (KRT76) Human Gene Symbol KRT77 , KRT76 Hu Chromosome Location 17q21.2 (KRT77) , 12q13.13 (KRT76) Synonym K1B , KRT1B , K77 , CK-1B , Keratin 1B , Keratin-77 , Cytokeratin-1B , KRT2B , KRT2P
Reactivity	Human, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG's
Clone Names	KRTL/1077 + KRTH/1076
Calculated MW	61901

Additional Information

Gene ID	374454
Other Names	Keratin, type II cytoskeletal 1b, Cytokeratin-1B, CK-1B, Keratin-77, K77, Type-II keratin Kb39, KRT77, KRT1B
Application Note	IF~~1:50~200 FC~~1:10~50 IHC-P~~N/A
Storage	Store at 2 to 8°C. Antibody is stable for 24 months.
Precautions	Cytokeratin, pan (Epithelial Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	KRT77
Synonyms	KRT1B
Tissue Location	Expressed exclusively in skin.

Background


Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0)

subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, which include CK1, CK3, CK4, CK5, CK6, CK8, CK10, CK14, CK15, CK16, and CK19. Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis. KRTL/KRTH is a broad spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It has been used to characterize the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms. This antibody stains cytokeratins present in normal and abnormal human tissues and has shown high sensitivity in the recognition of epithelial cells and carcinomas.


References

Woodcock-Mitchell J et. al. Journal of Cell Biology 1982;95:580-8.,2. Tseng SCG et. al. Cell 1982; 30361

Images



Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with Cytokeratin, pan Monoclonal Antibody cocktail (KRTL/1077 + KRTH/1076).



Formalin paraffin Rat Stomach stained with Cytokeratin, pan Monoclonal Antibody cocktail (KRTL/1077 + KRTH/1076).



Formalin-fixed, paraffin-embedded Rat Oviduct with Cytokeratin, pan Monoclonal Antibody cocktail (KRTL/1077 + KRTH/1076).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.