

Anti-CD257 / BAFF / TNFSF13B Antibody

Mouse Monoclonal Antibody

Catalog # AH13091

Product Information

Application	IF, FC
Primary Accession	Q9Y275
Other Accession	525157
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1
Clone Names	C257/1638
Calculated MW	31223

Additional Information

Gene ID	10673
Other Names	ApoL related ligand TALL 1; B lymphocyte stimulator; B-cell-activating factor; BAFF; BLyS; CD257; Dendritic cell-derived TNF-like molecule; DTL; PRO738; TALL-1; THANK; TNFSF13B; TNFSF20; Tumor necrosis factor ligand superfamily member 13b; Tumor necrosis factor ligand superfamily member 20; ZTNF4
Application Note	Flow Cytometry (0.5-1ug/million cells); ,Immunofluorescence (1-2ug/ml) ,Optimal dilution for a specific application should be determined.
Format	200ug/ml of Ab purified from rabbit anti-serum by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml.
Storage	Store at 2 to 8°C.Antibody is stable for 24 months.
Precautions	Anti-CD257 / BAFF / TNFSF13B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TNFSF13B
Synonyms	BAFF, BLyS, TALL1, TNFSF20, ZTNF4
Function	Cytokine that binds to TNFRSF13B/TACI and TNFRSF17/BCMA. TNFSF13/APRIL binds to the same 2 receptors. Together, they form a 2 ligands -2 receptors pathway involved in the stimulation of B- and T- cell function and the regulation of humoral immunity. A third B-cell specific BAFF-receptor

(BAFFR/BR3) promotes the survival of mature B- cells and the B-cell response.

Cellular Location

Cell membrane; Single-pass type II membrane protein

Tissue Location

Abundantly expressed in peripheral blood Leukocytes and is specifically expressed in monocytes and macrophages. Also found in the spleen, lymph node, bone marrow, T-cells and dendritic cells. A lower expression seen in placenta, heart, lung, fetal liver, thymus, and pancreas. Isoform 2 is expressed in many myeloid cell lines

Background

BAFF/CD257 is a type II transmembrane protein, and a member of the tumor necrosis factor ligand superfamily. It is proteolytically cleaved to form a soluble protein. BAFF, like other TNF ligand family members, forms homo-trimers. The predicted molecular weight of a BAFF monomer is approximately 31kDa. It is expressed on peripheral blood B and T lymphocytes, monocytes, macrophages, and dendritic cells and is upregulated by IFN- γ and down-regulated by PMA/ionomycin treatment. BAFF stimulates B and T cell immunity and regulates humoral immunity by binding TACI and BCMA receptors; BAFF prevents apoptosis and promotes B cell survival by binding the BAFF-specific receptor (BAFFR/BR3).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.