10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-Adipophilin / Perilipin-2 (Marker of Obesity) Antibody

Mouse Monoclonal Antibody
Catalog # AH13123

## Product Information

| | |
|---|---|
| **Application** | WB, IF, FC |
| **Primary Accession** | Q99541 |
| **Other Accession** | 3416 |
| **Reactivity** | Human |
| **Host** | Mouse |
| **Clonality** | Monoclonal |
| **Isotype** | Mouse / IgG1, kappa |
| **Clone Names** | ADFP/1365 |
| **Calculated MW** | 48075 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 123 |
| **Other Names** | Adipophilin; ADFP; Adipose differentiation-related protein (ADRP); Perilipin-2 (PLIN2) |
| **Application Note** | Flow Cytometry (0.5-1ug/million cells); ,Immunofluorescence (0.5-1ug/ml); ,Western Blotting (0.5-1ug/ml),Optimal dilution for a specific application should be determined. |
| **Format** | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| **Storage** | Store at 2 to 8°C.Antibody is stable for 24 months. |
| **Precautions** | Anti-Adipophilin / Perilipin-2 (Marker of Obesity) Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information


| | |
|---|---|
| **Name** | PLIN2 ( HGNC:248) |
| **Synonyms** | ADFP |
| **Function** | Structural component of lipid droplets, which is required for the formation and maintenance of lipid storage droplets.<br><br>Membrane {ECO:0000250\|UniProtKB:P43883}; Peripheral membrane protein |

| | |
|---|---|
| **Cellular Location** | {ECO:0000250\|UniProtKB:P43883}. Lipid droplet |
| **Tissue Location** | Milk lipid globules.. |

## Background

Recognizes a protein of 48kDa, which is identified as Adipophilin. It belongs to the perilipin family, members of which coat intracellular lipid storage droplets. This protein is associated with the lipid globule surface membrane material, and maybe involved in development and maintenance of adipose tissue. However, it is not restricted to adipocytes as previously thought, but is found in a wide range of cultured cell lines, including fibroblasts, endothelial and epithelial cells, and tissues, such as lactating mammary gland, adrenal cortex, Sertoli and Leydig cells, and hepatocytes in alcoholic liver cirrhosis, suggesting that it may serve as a marker of lipid accumulation in diverse cell types and diseases.

## Images




Western Blot of recombinant Adipophilin and Jurkat Cell Lysate using Adipophilin Monoclonal Antibody (ADFP/1365).



Flow Cytometry of human Adipophilin on PBMC. Green: Isotype Control; Red: Adipophilin Monoclonal Antibody (ADFP/1365).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.