

Anti-Growth Hormone Antibody

Mouse Monoclonal Antibody Catalog # AH13261

Product Information

| Application | IHC-P, IF, FC |
|-------------------|---------------|
| Primary Accession | <u>P01241</u> |
| Other Accession | <u>655229</u> |
| Reactivity | Human |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype | Mouse / IgG2b |
| Clone Names | GH/1450 |
| Calculated MW | 24847 |

Additional Information

| Gene ID | 2688 |
|------------------|---|
| Other Names | GH; GH-N; GH1; GHN; Growth hormone 1; Growth hormone; Growth hormone, pituitary; HG1; hGH-N; IGHD1B; Pituitary growth hormone; RNGHGP; Somatotropin |
| Application Note | Flow Cytometry (0.5-1ug/million cells); Immunofluorescence (1-2ug/ml); ,Immunohistology (Formalin-fixed) (0.5-1ug/ml for 30 min at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined. |
| Format | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Storage | Store at 2 to 8°C.Antibody is stable for 24 months. |
| Precautions | Anti-Growth Hormone Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| Name | GH1 |
|----------|---|
| Function | Plays an important role in growth control. Its major role in stimulating body growth is to stimulate the liver and other tissues to secrete IGF1. It stimulates both the differentiation and proliferation of myoblasts. It also stimulates amino acid uptake and protein synthesis in muscle and other tissues. |

Background

Pituitary growth hormone (GH) plays a crucial role in stimulating and controlling the growth, metabolism and differentiation of many mammalian cell types by modulating the synthesis of multiple mRNA species. These effects are mediated by the binding of GH to its membrane-bound receptor, GHR, and involve a phosphorylation cascade that results in the modulation of numerous signaling pathways. GH is synthesized by acidophilic or somatotropic cells of the anterior pituitary gland. Anti-GH is a useful marker in classification of pituitary tumors and the study of pituitary disease (acromegaly).

Images



Formalin-fixed, paraffin-embedded Human Pituitary stained with Growth Hormone Monoclonal Antibody (GH/1450).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.