

Anti-Connexin 32 (Gap Junction Protein) Antibody

Mouse Monoclonal Antibody

Catalog # AH13265

Product Information

Application	WB, IHC-F, IF, FC
Primary Accession	P08034
Other Accession	333303
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG
Clone Names	M12.13
Calculated MW	32025

Additional Information

Gene ID	2705
Other Names	Charcot Marie Tooth neuropathy X linked; CMTX; CMTX1; Connexin-32; Cx32; GAP junction 28kDa liver protein; Gap junction beta-1 protein; Gap junction protein beta 1 32kD; GJB1
Application Note	Immunofluorescence (1-2ug/ml); Flow Cytometry (0.5-1ug/million cells);,Western Blotting (0.5-1ug/ml); ,Immunohistology (Frozen) (0.5-1.0ug/ml for 30 minutes at RT),Optimal dilution for a specific application should be determined.
Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage	Store at 2 to 8°C.Antibody is stable for 24 months.
Precautions	Anti-Connexin 32 (Gap Junction Protein) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GJB1
Synonyms	CX32
Function	One gap junction consists of a cluster of closely packed pairs of transmembrane channels, the connexons, through which materials of low MW diffuse from one cell to a neighboring cell.

Cellular Location

Cell membrane; Multi-pass membrane protein. Cell junction, gap junction

Background

This Ab recognizes a protein of 27-32kDa, identified as Connexin 32. The connexin family of proteins forms hexameric complexes called connexons that facilitate movement of low molecular weight proteins between cells via gap junctions. Connexin proteins share a common topology of four transmembrane α -helical domains, two extracellular loops, a cytoplasmic loop and cytoplasmic N- and C-termini. Many of the key functional differences arise from specific amino-acid substitutions in the most highly conserved domains, the transmembrane and extracellular regions. Each of the approximately 20-connexin isoforms produces channels with distinct permeability and electrical and chemical sensitivities; therefore, one connexin usually cannot fully substitute for another.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.