10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Cytokeratin 10 Antibody

Mouse Monoclonal Antibody
Catalog # AH13353

Product Information

Application	IHC-P, IF, FC
Primary Accession	P13645
Other Accession	99936
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1 s, kappa
Clone Names	KRT10/844 + KRT10/1275
Calculated MW	58827

Additional Information

Gene ID	3858
Other Names	BCIE, BIE, EHK, Keratin Type I Cytoskeletal 10, KRT10
Application Note	Flow Cytometry (0.5-1ug/million cells in 0.1ml); Immunofluorescence (0.5-1ug/ml); ,Immunohistology (Formalin-fixed) (0.1-0.2ug/ml for 30 min at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.
Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage	Store at 2 to 8°C.Antibody is stable for 24 months.
Precautions	Anti-Cytokeratin 10 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	KRT10 (HGNC:6413)
Synonyms	KPP
Function	Structural component of intermediate filaments in suprabasal keratinocytes of stratified epithelia. Forms heteropolymers primarily with the type II keratin KRT1, and also with KRT2, assembling into intermediate filament networks that support terminal differentiation, mechanical resilience, and epidermal

	barrier formation.
Cellular Location	Secreted, extracellular space. Cell surface. Cytoplasm
Tissue Location	Seen in all suprabasal cell layers including stratum corneum. Expressed on the surface of lung cell lines (PubMed:19627498). Localized on the surface of desquamated nasal epithelial cells (at protein level) (PubMed:12427098)

Background

This MAb recognizes a protein of 56.5kDa, identified as cytokeratin 10 (CK10). CK10 is expressed in all suprabasal layers of the epidermis. In the epidermis, expression of CK10 strictly parallels the extent of differentiation; it is absent in the basal layer, appears in the first suprabasal layers and increases in concentration towards the granular layer. However, CK10 is rarely detected in early stages of vulvar squamous carcinomas (tumors less than 2 cm, clinical stage I) regardless of the tumor grade. In larger and more advanced tumors (greater than 2 cm, clinical stages II and III), CK10 is detected very frequently. Expression of CK10 is related to maturation of malignant keratinocytes, being preferentially detected in more-differentiated parts.

Images



Formalin-fixed, paraffin-embedded human Skin stained with Cytokeratin 10 Monoclonal Antibody (KRT10/+ KRT10/1275).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.