

Anti-NKX2.2 Antibody

Mouse Monoclonal Antibody

Catalog # AH13411

Product Information

Application	IHC-P, IF, FC
Primary Accession	O95096
Other Accession	516922
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Clone Names	NX2/1523
Calculated MW	30133

Additional Information

Gene ID	4821
Other Names	Homeobox protein NK-2 homolog B, NK2 transcription factor like protein B, NK2 transcription factor related locus 2, NKX22, Nkx2b, tinman
Application Note	Flow Cytometry (0.5-1ug/million cells); Immunofluorescence (1-2ug/ml); ,Immunohistology (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.
Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage	Store at 2 to 8°C.Antibody is stable for 24 months.
Precautions	Anti-NKX2.2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NKX2-2
Synonyms	NKX2.2, NKX2B
Function	Transcriptional activator involved in the development of insulin-producing beta cells in the endocrine pancreas (By similarity). May also be involved in specifying diencephalic neuromeric boundaries, and in controlling the

expression of genes that play a role in axonal guidance. Binds to elements within the NEUROD1 promoter (By similarity).


Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108}.


Background

Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein specific to Ewing sarcoma, and was shown to be differentially upregulated in Ewing sarcoma on the basis of array-based gene expression analysis. It acts as a valuable marker for Ewing sarcoma, with a sensitivity of 93% and a specificity of 89%, and aids in the differential diagnosis of small round cell tumors.

Images



Formalin-fixed, paraffin-embedded human Ewing Sarcoma stained with NKX2.2 Monoclonal Antibody (NX2/1523).



Formalin-fixed, paraffin-embedded human Ewing Sarcoma stained with NKX2.2 Monoclonal Antibody (NX2/1523).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.