

# Anti-Wilm s Tumor 1 (WT1) Antibody

Recombinant Mouse Monoclonal Antibody

Catalog # AH13578

## Product Information

---

Application	IHC-P, IF, FC
Primary Accession	<a href="#">P19544</a>
Other Accession	<a href="#">591980</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Clone Names	rWT1/857
Calculated MW	49188

## Additional Information

---

Gene ID	7490
Other Names	WT1; AWT1; FWT1; GUD; NPHS4; WAGR; Wilms tumor 1
Application Note	Flow Cytometry (0.5-1ug/million cells); Immunofluorescence (0.5-1ug/ml); ,Immunohistology (Formalin-fixed) (0.5-1ug/ml for 30 minutes at RT),Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.
Format	200ug/ml of recombinant MAb purified by Protein A/G. Prepared in 1mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage	Store at 2 to 8°C.Antibody is stable for 24 months.
Precautions	Anti-Wilm s Tumor 1 (WT1) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	WT1
Function	Transcription factor that plays an important role in cellular development and cell survival (PubMed: <a href="#">7862533</a> ). Recognizes and binds to the DNA sequence 5'-GCG(T/G)GGGCG-3' (PubMed: <a href="#">17716689</a> , PubMed: <a href="#">25258363</a> , PubMed: <a href="#">7862533</a> ). Regulates the expression of numerous target genes, including EPO. Plays an essential role for development of the urogenital system. It has a tumor suppressor as well as an oncogenic role in tumor

formation. Function may be isoform-specific: isoforms lacking the KTS motif may act as transcription factors (PubMed:[15520190](#)). Isoforms containing the KTS motif may bind mRNA and play a role in mRNA metabolism or splicing (PubMed:[16934801](#)). Isoform 1 has lower affinity for DNA, and can bind RNA (PubMed:[19123921](#)).

#### Cellular Location

Nucleus. Nucleus, nucleolus. Cytoplasm. Note=Isoforms lacking the KTS motif have a diffuse nuclear location (PubMed:15520190). Shuttles between nucleus and cytoplasm. {ECO:0000250, ECO:0000269 | PubMed:15520190} [Isoform 4]: Nucleus, nucleoplasm

#### Tissue Location

Expressed in the kidney and a subset of hematopoietic cells


## Background

---

Recognizes a 47-55kDa-tumor suppressor protein, identified as Wilm's Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subsets of sporadic Wilms tumors. WT1, a sporadic and familial pediatric kidney tumor, is genetically heterogeneous. Wilms tumor is associated with mutations of WT1, a zinc-finger transcription factor that is essential for the development of the metanephric kidney and the urogenital system. The WT1 gene is normally expressed in fetal kidney and mesothelium, and its expression has been suggested as a marker for Wilms tumor and mesothelioma. WT1 protein has been identified in proliferative mesothelial cells, malignant mesothelioma, ovarian carcinoma, gonadoblastoma, nephroblastoma, and desmoplastic small round cell tumor. Lung adenocarcinomas rarely stain positive with this antibody. WT1 protein expression in mesothelial cells has become a reliable marker for the diagnosis of mesotheliomas.

## Images

---



Formalin-fixed, paraffin-embedded Human Kidney stained with Wilm's Tumor Recombinant Mouse Monoclonal Antibody (rWT1/857).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.