

## **PPIA Antibody**

Mouse Monoclonal Antibody Catalog # AI10002

## **Product Information**

Application WB Primary Accession P62937

Other Accession P62937, NP 066953, NM 021130

Reactivity Human
Host Mouse
Clonality Monoclonal
Isotype Mouse IgG1, k

Clone Names 9G7 Calculated MW 18012

## **Additional Information**

**Gene ID** 5478

Alias Symbol CYPA, CYPH, MGC12404, MGC23397, MGC117158

Other Names Peptidyl-prolyl cis-trans isomerase A, PPIase A, Cyclophilin A, Cyclosporin

A-binding protein, Rotamase A, Peptidyl-prolyl cis-trans isomerase A,

N-terminally processed, PPIA, CYPA

**Target/Specificity** This gene encodes a member of the peptidyl-prolyl cis-trans isomerase

(PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. The encoded protein is a cyclosporin binding-protein and may play a role in cyclosporin A-mediated immunosuppression. The protein can also interact with several HIV proteins, including p55 gag, Vpr, and capsid protein, and has been shown to be necessary for the formation of infectious HIV virions. Multiple pseudogenes that map to different chromosomes have been

reported.

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 100ul of distilled water. Final anti-PPIA antibody concentration is 1 mg/ml

in PBS buffer. For longer periods of storage, store at -20°C. Avoid repeat

freeze-thaw cycles.

**Precautions** PPIA Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

## **Protein Information**

Name PPIA

**Synonyms** CYPA

**Function** 

Catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides (PubMed:2001362, PubMed:20676357, PubMed:21245143, PubMed: 21593166, PubMed: 25678563). Exerts a strong chemotactic effect on leukocytes partly through activation of one of its membrane receptors BSG/CD147, initiating a signaling cascade that culminates in MAPK/ERK activation (PubMed: 11943775, PubMed: 21245143). Activates endothelial cells (ECs) in a pro-inflammatory manner by stimulating activation of NF-kappa-B and ERK, INK and p38 MAP-kinases and by inducing expression of adhesion molecules including SELE and VCAM1 (PubMed:15130913). Induces apoptosis in ECs by promoting the FOXO1-dependent expression of CCL2 and BCL2L11 which are involved in EC chemotaxis and apoptosis (PubMed:31063815). In response to oxidative stress, initiates proapoptotic and antiapoptotic signaling in ECs via activation of NF-kappa-B and AKT1 and up-regulation of antiapoptotic protein BCL2 (PubMed:23180369). Negatively regulates MAP3K5/ASK1 kinase activity, autophosphorylation and oxidative stress-induced apoptosis mediated by MAP3K5/ASK1 (PubMed:26095851). Necessary for the assembly of TARDBP in heterogeneous nuclear ribonucleoprotein (hnRNP) complexes and regulates TARDBP binding to RNA UG repeats and TARDBP-dependent expression of HDAC6, ATG7 and VCP which are involved in clearance of protein aggregates (PubMed: 25678563). Plays an important role in platelet activation and aggregation (By similarity). Regulates calcium mobilization and integrin ITGA2B:ITGB3 bidirectional signaling via increased ROS production as well as by facilitating the interaction between integrin and the cell cytoskeleton (By similarity). Binds heparan sulfate glycosaminoglycans (PubMed:11943775). Inhibits replication of influenza A virus (IAV) (PubMed: 19207730). Inhibits ITCH/AIP4-mediated ubiquitination of matrix protein 1 (M1) of IAV by impairing the interaction of ITCH/AIP4 with M1, followed by the suppression of the nuclear export of M1, and finally reduction of the replication of IAV (PubMed: 22347431, PubMed:30328013).

**Cellular Location** 

Cytoplasm. Secreted. Nucleus Note=Secretion occurs in response to oxidative stress in vascular smooth muscle through a vesicular secretory pathway that includes Rho GTPase signaling, actin remodeling, and myosin II activation

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.