

LASS2 antibody - N-terminal region

Rabbit Polyclonal Antibody Catalog # AI10036

Product Information

Application WB Primary Accession Q96G23

Other Accession <u>096G23</u>, <u>CAI13332</u>, <u>NM 022075</u>

Reactivity Human, Mouse, Rat, Rabbit, Dog, Guinea Pig, Horse, Bovine

Predicted Mouse, Pig, Horse

Host Rabbit
Clonality Polyclonal
Calculated MW 44876

Additional Information

Gene ID 29956

Alias Symbol CerS2, FLJ10243, L3, MGC987, SP260, TMSG1, LASS2

Other Names Ceramide synthase 2, CerS2, LAG1 longevity assurance homolog 2, SP260,

Tumor metastasis-suppressor gene 1 protein, CERS2, LASS2, TMSG1

Target/Specificity LASS2 is a protein that has sequence similarity to yeast longevity assurance

gene 1. Mutation or overexpression of the related gene in yeast has been shown to alter yeast lifespan. The human protein may play a role in the regulation of cell growth. This gene encodes a protein that has sequence similarity to yeast longevity assurance gene 1. Mutation or overexpression of the related gene in yeast has been shown to alter yeast lifespan. The human protein may play a role in the regulation of cell growth. Alternatively spliced

transcript variants encoding the same protein have been described.

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 50 ul of distilled water. Final anti-LASS2 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

-20°C. Avoid repeat freeze-thaw cycles.

Precautions LASS2 antibody - N-terminal region is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name CERS2 {ECO:0000303|PubMed:17977534, ECO:0000312|HGNC:HGNC:14076}

Function Ceramide synthase that catalyzes the transfer of the acyl chain from

acyl-CoA to a sphingoid base, with high selectivity toward very-long-chain fatty acyl-CoA (chain length C22-C27) (PubMed: 17977534, PubMed: 18165233,

PubMed: 18541923, PubMed: 19728861, PubMed: 20937905,

PubMed:22144673, PubMed:22661289, PubMed:26887952,

PubMed:<u>29632068</u>). N- acylates sphinganine and sphingosine bases to form dihydroceramides and ceramides in de novo synthesis and salvage pathways, respectively (By similarity) (PubMed:<u>17977534</u>, PubMed:<u>18165233</u>,

PubMed:<u>18541923</u>, PubMed:<u>19728861</u>, PubMed:<u>20937905</u>, PubMed:<u>22144673</u>, PubMed:<u>22661289</u>, PubMed:<u>26887952</u>,

PubMed: 29632068). Plays a non-redundant role in the synthesis of ceramides with very-long-chain fatty acids in kidney, liver and brain. Regulates the abundance of myelin-specific sphingolipids galactosylceramide and sulfatide that affects myelin sheath architecture and motor neuron functions (By

similarity).


Cellular Location Endoplasmic reticulum membrane; Multi-pass membrane protein

Tissue Location Expressed in kidney, liver, brain, heart, placenta and lung.

Background

This is a rabbit polyclonal antibody against LASS2. It was validated on Western Blot using a cell lysate as a positive control. Abgent strives to provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).

Images



LASS2 antibody - N-terminal region (AI10036) in Human HepG2 cells using Western Blot WB Suggested Anti-LASS2 Antibody Titration: 0.2-1 μ g/ml Positive Control: HepG2 cell lysate CERS2 is supported by BioGPS gene expression data to be expressed in HepG2

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.