

CNOT7 antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI10068

Product Information

Application	WB
Primary Accession	Q9UIV1
Other Accession	Q9UIV1 , NP_037486 , NM_013354
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Goat, Dog, Guinea Pig, Horse, Bovine, Yeast
Predicted Host	Human, Mouse, Rat, Zebrafish, Pig, Chicken, Goat, Dog, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 32745

Additional Information

Gene ID	29883
Alias Symbol	CAF1, hCAF-1
Other Names	CCR4-NOT transcription complex subunit 7, BTG1-binding factor 1, CCR4-associated factor 1, CAF-1, Caf1a, CNOT7, CAF1
Target/Specificity	CNOT7 binds to an anti-proliferative protein, B-cell translocation protein 1, which negatively regulates cell proliferation. Binding of the two proteins, which is driven by phosphorylation of the anti-proliferative protein, causes signaling events in cell division that lead to changes in cell proliferation associated with cell-cell contact. The protein has both mouse and yeast orthologs.
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-CNOT7 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.
Precautions	CNOT7 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CNOT7
Synonyms	CAF1


Function Has 3'-5' poly(A) exoribonuclease activity for synthetic poly(A) RNA substrate (PubMed:[19276069](#), PubMed:[20634287](#), PubMed:[31439799](#)). Its function seems to be partially redundant with that of CNOT8 (PubMed:[19605561](#)). Catalytic component of the CCR4-NOT complex which is one of the major cellular mRNA deadenylases and is linked to various cellular processes including bulk mRNA degradation, miRNA-mediated repression, translational repression during translational initiation and general transcription regulation (PubMed:[19276069](#), PubMed:[20634287](#), PubMed:[31439799](#)). During miRNA-mediated repression the complex also seems to act as translational repressor during translational initiation (PubMed:[20065043](#)). Additional complex functions may be a consequence of its influence on mRNA expression (PubMed:[19276069](#), PubMed:[23236473](#)). Associates with members of the BTG family such as TOB1 and BTG2 and is required for their anti-proliferative activity (PubMed:[19276069](#), PubMed:[23236473](#)).

Cellular Location Nucleus. Cytoplasm, P-body {ECO:0000250|UniProtKB:Q60809}. Cytoplasm, Cytoplasmic ribonucleoprotein granule. Note=NANOS2 promotes its localization to P-body (By similarity). Recruited to cytoplasmic ribonucleoprotein membraneless compartments by CAPRIN1, promoting deadenylation of mRNAs (PubMed:31439799) {ECO:0000250|UniProtKB:Q60809, ECO:0000269|PubMed:31439799}

Background

This is a rabbit polyclonal antibody against CNOT7. It was validated on Western Blot using a cell lysate as a positive control. Abgent strives to provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).

Images



CNOT7 antibody - middle region (AI10068) in Human Placenta cells using Western Blot
WB Suggested Anti-CNOT7 Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:62500
Positive Control: Human Placenta

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.