

BRD3 Antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI10087

Product Information

Application	WB
Primary Accession	Q15059
Other Accession	Q15059-2
Reactivity	Human
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	79542

Additional Information

Gene ID	8019
Other Names	Bromodomain-containing protein 3, RING3-like protein, BRD3, KIAA0043, RING3L
Target/Specificity	This gene was identified based on its homology to the gene encoding the RING3 protein, a serine/threonine kinase. The gene localizes to 9q34, a region which contains several major histocompatibility complex (MHC) genes. The function of the encoded protein is not known.
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul, l of distilled water. Final Anti-BRD3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.
Precautions	BRD3 Antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	BRD3 {ECO:0000303 PubMed:18406326, ECO:0000312 HGNC:HGNC:1104}
Function	Chromatin reader that recognizes and binds acetylated histones, thereby controlling gene expression and remodeling chromatin structures (PubMed: 18406326 , PubMed: 22464331 , PubMed: 27105114 , PubMed: 32895492). Recruits transcription factors and coactivators to target gene sites, and activates RNA polymerase II machinery for transcriptional elongation (PubMed: 29567837 , PubMed: 32895492). In vitro, binds acetylated

lysine residues on the N-terminus of histone H2A, H2B, H3 and H4 (PubMed:[18406326](#)). Involved in endoderm differentiation via its association with long non-coding RNA (lncRNA) DIGIT: BRD3 undergoes liquid-liquid phase separation upon binding to lncRNA DIGIT, promoting binding to histone H3 acetylated at 'Lys-18' (H3K18ac) to induce endoderm gene expression (PubMed:[32895492](#)). Also binds non-histones acetylated proteins, such as GATA1 and GATA2: regulates transcription by promoting the binding of the transcription factor GATA1 to its targets (By similarity).

Cellular Location

Nucleus. Chromosome. Note=Detected on chromatin

Tissue Location

Ubiquitous..

Background

This is a rabbit polyclonal antibody against BRD3. It was validated on Western Blot by Abgent. At Abgent we manufacture rabbit polyclonal antibodies on a large scale (200-1000 products/month) of high throughput manner. Our antibodies are peptide based and protein family oriented. We usually provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).

Images



BRD3 Antibody - middle region (AI10087) in Human Liver tumor cells using Western Blot

Host: Rabbit

Target Name: BRD3

Sample Tissue: Liver tumor lysates

Antibody Dilution: 1.0 µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.