

PARP2 Antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI10104

Product Information

Application	WB
Primary Accession	Q9UGN5
Other Accession	Q9UGN5-2 , NP_001036083 , NM_001042618
Reactivity	Human, Mouse, Rat, Rabbit, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Pig, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	66206

Additional Information

Gene ID	10038
Alias Symbol Other Names	ADPRT2, ADPRTL2, ADPRTL3, ARTD2, PARP-2, pADPRT-2 Poly [ADP-ribose] polymerase 2, PARP-2, hPARP-2, ADP-ribosyltransferase diphtheria toxin-like 2, ARTD2, NAD(+) ADP-ribosyltransferase 2, ADPRT-2, Poly[ADP-ribose] synthase 2, pADPRT-2, PARP2, ADPRT2, ADPRTL2
Target/Specificity	PARP2 is involved in the base excision repair (BER) pathway, by catalyzing the poly(ADP-ribosyl)ation of a limited number of acceptor proteins involved in chromatin architecture and in DNA metabolism. This modification follows DNA damages and appears as an obligatory step in a detection/signaling pathway leading to the reparation of DNA strand breaks.
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul, l of distilled water. Final Anti-PARP2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.
Precautions	PARP2 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PARP2 {ECO:0000303 PubMed:20092359, ECO:0000312 HGNC:HGNC:272}
Function	Poly-ADP-ribosyltransferase that mediates poly-ADP- ribosylation of proteins and plays a key role in DNA repair (PubMed: 10364231 , PubMed: 25043379 , PubMed: 27471034 , PubMed: 30104678 , PubMed: 32028527 ,

PubMed:[32939087](#), PubMed:[34108479](#), PubMed:[34486521](#), PubMed:[34874266](#)). Mediates glutamate, aspartate or serine ADP-ribosylation of proteins: the ADP-D-ribosyl group of NAD(+) is transferred to the acceptor carboxyl group of target residues and further ADP-ribosyl groups are transferred to the 2'-position of the terminal adenosine moiety, building up a polymer with an average chain length of 20-30 units (PubMed:[25043379](#), PubMed:[30104678](#), PubMed:[30321391](#)). Serine ADP-ribosylation of proteins constitutes the primary form of ADP-ribosylation of proteins in response to DNA damage (PubMed:[32939087](#)). Mediates glutamate and aspartate ADP-ribosylation of target proteins in absence of HPF1 (PubMed:[25043379](#)). Following interaction with HPF1, catalyzes serine ADP-ribosylation of target proteins; HPF1 conferring serine specificity by completing the PARP2 active site (PubMed:[28190768](#), PubMed:[32028527](#), PubMed:[34108479](#), PubMed:[34486521](#), PubMed:[34874266](#)). PARP2 initiates the repair of double-strand DNA breaks: recognizes and binds DNA breaks within chromatin and recruits HPF1, licensing serine ADP-ribosylation of target proteins, such as histones, thereby promoting decompaction of chromatin and the recruitment of repair factors leading to the reparation of DNA strand breaks (PubMed:[10364231](#), PubMed:[32939087](#), PubMed:[34108479](#)). HPF1 initiates serine ADP-ribosylation but restricts the polymerase activity of PARP2 in order to limit the length of poly-ADP-ribose chains (PubMed:[34732825](#), PubMed:[34795260](#)). Specifically mediates formation of branched poly-ADP-ribosylation (PubMed:[30104678](#)). Branched poly-ADP-ribose chains are specifically recognized by some factors, such as APLF (PubMed:[30104678](#)). In addition to proteins, also able to ADP-ribosylate DNA: preferentially acts on 5'-terminal phosphates at DNA strand breaks termini in nicked duplex (PubMed:[27471034](#), PubMed:[29361132](#)).

Cellular Location

Nucleus. Chromosome. Note=Recruited to DNA damage sites in a PARP1-dependent process: recognizes and binds poly-ADP-ribose chains produced by PARP1 at DNA damage sites via its N-terminus, leading to its recruitment.


Tissue Location

Widely expressed, mainly in actively dividing tissues (PubMed:[10364231](#)). The highest levels are in the brain, heart, pancreas, skeletal muscle and testis; also detected in kidney, liver, lung, placenta, ovary and spleen; levels are low in leukocytes, colon, small intestine, prostate and thymus (PubMed:[10364231](#))

Background

This is a rabbit polyclonal antibody against PARP2. It was validated on Western Blot by Abgent. At Abgent we manufacture rabbit polyclonal antibodies on a large scale (200-1000 products/month) of high throughput manner. Our antibodies are peptide based and protein family oriented. We usually provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).

Images



PARP2 Antibody - C-terminal region (AI10104) in Human MCF7 cells using Western Blot
Host: Rabbit
Target Name: PARP2
Sample Tissue: MCF7 Whole cell lysate
Antibody Dilution: 1.0 µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.