

TRIM9 antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI10121

Product Information

Application	WB
Primary Accession	Q9C026
Other Accession	Q9C026 , NP_055978 , NM_015163
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	79177

Additional Information

Gene ID	114088
Alias Symbol	KIAA0282, RNF91, SPRING
Other Names	E3 ubiquitin-protein ligase TRIM9, 632-, RING finger protein 91, Tripartite motif-containing protein 9, TRIM9, KIAA0282, RNF91
Target/Specificity	TRIM9 is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to cytoplasmic bodies. Its function has not been identified. Alternate splicing of this gene generates two transcript variants encoding different isoforms. The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to cytoplasmic bodies. Its function has not been identified. Alternate splicing of this gene generates two transcript variants encoding different isoforms.
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-TRIM9 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.
Precautions	TRIM9 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	TRIM9
-------------	-------

Synonyms	KIAA0282, RNF91
Function	E3 ubiquitin-protein ligase which ubiquitinates itself in cooperation with an E2 enzyme UBE2D2/UBC4 and serves as a targeting signal for proteasomal degradation. May play a role in regulation of neuronal functions and may also participate in the formation or breakdown of abnormal inclusions in neurodegenerative disorders. May act as a regulator of synaptic vesicle exocytosis by controlling the availability of SNAP25 for the SNARE complex formation.
Cellular Location	Cytoplasm. Cell projection, dendrite. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle {ECO:0000250 UniProtKB:Q91ZY8}. Synapse {ECO:0000250 UniProtKB:Q91ZY8} Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:Q91ZY8}. Note=Enriched at synaptic terminals where it exists in a soluble form and a synaptic vesicle-associated form. Associated with the cytoskeleton (By similarity). Found in proximal dendrites of pyramidal neurons in the cerebral cortex and hippocampus, and Purkinje cells in the cerebellum (PubMed:20085810). {ECO:0000250 UniProtKB:Q91ZY8, ECO:0000269 PubMed:20085810}
Tissue Location	Brain. Highly expressed in the cerebral cortex (at protein level). Severely decreased in the affected brain areas in Parkinson disease and dementia with Lewy bodies

Background

This is a rabbit polyclonal antibody against TRIM9. It was validated on Western Blot using a cell lysate as a positive control. Abgent strives to provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).

Images



TRIM9 antibody - middle region (AI10121) in Transfected 293T cells using Western Blot
 WB Suggested Anti-TRIM9 Antibody Titration: 0.2-1 µg/ml
 ELISA Titer: 1:62500
 Positive Control: Transfected 293T

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.