

# SART3 antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI10125

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">Q15020</a>
<b>Other Accession</b>	<a href="#">Q15020</a> , <a href="#">NP_055521</a> , <a href="#">NM_014706</a>
<b>Reactivity</b>	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine, Yeast
<b>Predicted</b>	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Chicken, Dog, Guinea Pig, Horse, Bovine
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	109935

## Additional Information

---

<b>Gene ID</b>	9733
<b>Alias Symbol</b> <b>Other Names</b>	KIAA0156, MGC138188, RP11-13G14, TIP110, p110(nrb), P100, p110, DSAP1 Squamous cell carcinoma antigen recognized by T-cells 3, SART-3 {ECO:0000312 EMBL:BAA783841}, Tat-interacting protein of 110 kDa, Tip110, p110 nuclear RNA-binding protein, SART3 ( <a href="#">HGNC:16860</a> )
<b>Target/Specificity</b>	<p>SART3 is an RNA-binding nuclear protein that is a tumor-rejection antigen. This antigen possesses tumor epitopes capable of inducing HLA-A24-restricted and tumor-specific cytotoxic T lymphocytes in cancer patients and may be useful for specific immunotherapy. SART3 is found to be an important cellular factor for HIV-1 gene expression and viral replication. It also associates transiently with U6 and U4/U6 snRNPs during the recycling phase of the spliceosome cycle. This protein is thought to be involved in the regulation of mRNA splicing. The protein encoded by this gene is an RNA-binding nuclear protein that is a tumor-rejection antigen. This antigen possesses tumor epitopes capable of inducing HLA-A24-restricted and tumor-specific cytotoxic T lymphocytes in cancer patients and may be useful for specific immunotherapy. This gene product is found to be an important cellular factor for HIV-1 gene expression and viral replication. It also associates transiently with U6 and U4/U6 snRNPs during the recycling phase of the spliceosome cycle. This encoded protein is thought to be involved in the regulation of mRNA splicing. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.</p>
<b>Format</b>	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
<b>Reconstitution &amp; Storage</b>	Add 50 ul of distilled water. Final anti-SART3 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

## Precautions

SART3 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.


## Protein Information

<b>Name</b>	SART3 ( <a href="#">HGNC:16860</a> )
<b>Function</b>	U6 snRNP-binding protein that functions as a recycling factor of the splicing machinery. Promotes the initial reassembly of U4 and U6 snRNPs following their ejection from the spliceosome during its maturation (PubMed: <a href="#">12032085</a> ). Also binds U6atac snRNPs and may function as a recycling factor for U4atac/U6atac spliceosomal snRNP, an initial step in the assembly of U12-type spliceosomal complex. The U12-type spliceosomal complex plays a role in the splicing of introns with non- canonical splice sites (PubMed: <a href="#">14749385</a> ). May also function as a substrate-targeting factor for deubiquitinases like USP4 and USP15. Recruits USP4 to ubiquitinated PRPF3 within the U4/U5/U6 tri-snRNP complex, promoting PRPF3 deubiquitination and thereby regulating the spliceosome U4/U5/U6 tri-snRNP spliceosomal complex disassembly (PubMed: <a href="#">20595234</a> ). May also recruit the deubiquitinase USP15 to histone H2B and mediate histone deubiquitination, thereby regulating gene expression and/or DNA repair (PubMed: <a href="#">24526689</a> ). May play a role in hematopoiesis probably through transcription regulation of specific genes including MYC (By similarity).
<b>Cellular Location</b>	Nucleus, nucleoplasm. Nucleus, Cajal body. Nucleus speckle. Cytoplasm
<b>Tissue Location</b>	Ubiquitously expressed.

## Background

This is a rabbit polyclonal antibody against SART3. It was validated on Western Blot using a cell lysate as a positive control. Abgent strives to provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire ([sales@abgent.com](mailto:sales@abgent.com)).

## Images



SART3 antibody - middle region (AI10125) in Human Small Intestine cells using Western Blot  
WB Suggested Anti-SART3 Antibody Titration: 0.2-1 µg/ml  
ELISA Titer: 1:62500  
Positive Control: Human Small Intestine

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.