

# **TNFRSF10B Antibody**

Mouse Antibody Catalog # AI10155

### **Product Information**

Application WB Primary Accession 014763

Other Accession <u>NM 003842, NP 671716</u>

Reactivity Human Host Mouse

**Isotype** Mouse IgM,k

Clone Names 11G7 Calculated MW 47878

## **Additional Information**

**Gene ID** 8795

Alias Symbol DR5, CD262, KILLER, TRICK2, TRICKB, ZTNFR9, TRAILR2, TRICK2A, TRICK2B,

TRAIL-R2, KILLER/DR5

Other Names Tumor necrosis factor receptor superfamily member 10B, Death receptor 5,

TNF-related apoptosis-inducing ligand receptor 2, TRAIL receptor 2, TRAIL-R2,

CD262, TNFRSF10B, DR5, KILLER, TRAILR2, TRICK2, ZTNFR9

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 100ul of distilled water. Final anti-TNFRSF10B antibody concentration is 1

mg/ml in PBS buffer. For longer periods of storage, store at -200°C. Avoid

repeat freeze-thaw cycles.

**Precautions**TNFRSF10B Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

## **Protein Information**

Name TNFRSF10B

**Synonyms** DR5, KILLER, TRAILR2, TRICK2, ZTNFR9

**Function** Receptor for the cytotoxic ligand TNFSF10/TRAIL (PubMed: 10549288). The

adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa-B. Essential for ER stress-induced apoptosis.


#### **Cellular Location**

Membrane; Single-pass type I membrane protein.

#### **Tissue Location**

Widely expressed in adult and fetal tissues; very highly expressed in tumor cell lines such as HeLaS3, K-562, HL-60, SW480, A-549 and G-361; highly expressed in heart, peripheral blood lymphocytes, liver, pancreas, spleen, thymus, prostate, ovary, uterus, placenta, testis, esophagus, stomach and throughout the intestinal tract; not detectable in brain

## **Images**



WB Suggested Anti-TNFRSF1B Antibody Titration: 1 μg/ml Positive Control: Placenta

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.