

Wnt2 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI10314

Product Information


WB
<u>P21552</u>
<u>NM_023653</u> , <u>NP_076142</u>
Human, Mouse, Rat, Zebrafish, Pig, Dog, Bovine
Human, Mouse, Rat, Zebrafish, Pig, Dog, Bovine
Rabbit
Polyclonal
40482

Additional Information

Gene ID	22413
Alias Symbol Other Names	2610510E18Rik, Int1l1, Irp, Mirp, Wnt-2, Wnt2a Protein Wnt-2, INT-1-related protein, IRP, Wnt2, Irp, Wnt-2
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Wnt2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Wnt2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Wnt2
Synonyms	Irp, Wnt-2
Function	Ligand for members of the frizzled family of seven transmembrane receptors. Functions in the canonical Wnt/beta-catenin signaling pathway (PubMed: <u>19686689</u>). Functions as a upstream regulator of FGF10 expression (PubMed: <u>19686689</u>). Plays an important role in embryonic lung development (PubMed: <u>19686689</u>). May contribute to embryonic brain development by regulating the proliferation of dopaminergic precursors and neurons (PubMed: <u>20018874</u>).
Cellular Location	Secreted, extracellular space, extracellular matrix



Citations

29 kDa_ 22 kDa_

• Overexpression of miR I214 promotes the progression of human osteosarcoma by regulating the Wnt/β Icatenin signaling pathway.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.