

GLI1 antibody - C-terminal region

Rabbit Polyclonal Antibody
Catalog # AI10350

Product Information

Application	WB
Primary Accession	P08151
Other Accession	NM_005269 , NP_005260
Reactivity	Human, Mouse, Rat, Pig, Dog, Bovine
Predicted	Human, Mouse, Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	117904

Additional Information

Gene ID	2735
Alias Symbol	GLI
Other Names	Zinc finger protein GLI1, Glioma-associated oncogene, Oncogene GLI, GLI1, GLI
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 100 ul of distilled water. Final anti-GLI1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	GLI1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GLI1
Synonyms	GLI
Function	Acts as a transcriptional activator (PubMed: 10806483 , PubMed: 19706761 , PubMed: 19878745 , PubMed: 24076122 , PubMed: 24217340 , PubMed: 24311597). Binds to the DNA consensus sequence 5'-GACCACCCA-3' (PubMed: 2105456 , PubMed: 24217340 , PubMed: 8378770). Regulates the transcription of specific genes during normal development (PubMed: 19706761). Plays a role in craniofacial development and digital development, as well as development of the central nervous system and gastrointestinal tract. Mediates SHH signaling (PubMed: 19706761 ,

PubMed:[28973407](#)). Plays a role in cell proliferation and differentiation via its role in SHH signaling (PubMed:[11238441](#), PubMed:[28973407](#)).

Cellular Location

Cytoplasm. Nucleus. Note=Tethered in the cytoplasm by binding to SUFU (PubMed:10806483). Activation and translocation to the nucleus is promoted by interaction with STK36 (PubMed:10806483). Phosphorylation by ULK3 may promote nuclear localization (PubMed:19878745). Translocation to the nucleus is promoted by interaction with ZIC1 (PubMed:11238441)


Tissue Location

Detected in testis (at protein level) (PubMed:2105456). Testis, myometrium and fallopian tube. Also expressed in the brain with highest expression in the cerebellum, optic nerve and olfactory tract (PubMed:19878745). Isoform 1 is detected in brain, spleen, pancreas, liver, kidney and placenta; isoform 2 is not detectable in these tissues (PubMed:19706761)

References


Rahnama, F., et al., (2006) *Biochem. J.* 394 (PT 1), 19-26
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.
Publications: Yoshimura, K., Kawate, T. & Takeda, S. Signaling through the primary cilium affects glial cell survival under a stressed environment. *Glia* 59, 333-44 (2011). WB, IP, Bovine, Human, Dog, Pig, Horse, Rabbit, Rat, Guinea pig, Mouse21125655
Maitah, M. Y., Ali, S., Ahmad, A., Gadgeel, S. & Sarkar, F. H. Up-regulation of sonic hedgehog contributes to TGF- β 1-induced epithelial to mesenchymal transition in NSCLC cells. *PLoS One* 6, e16068 (2011). WB, IP, Bovine, Human, Dog, Pig, Horse, Rabbit, Rat, Guinea pig, Mouse21249152
Schiapparelli, P. et al. Inhibition of the sonic hedgehog pathway by cyclopamine reduces the CD133+/CD15+ cell compartment and the in vitro tumorigenic capability of neuroblastoma cells. *Cancer Lett.* 310, 222-31 (2011). WB, IP, Bovine, Human, Dog, Pig, Horse, Rabbit, Rat, Guinea pig, Mouse21803487
Ferruzzi, P. et al. In vitro and in vivo characterization of a novel Hedgehog signaling antagonist in human glioblastoma cell lines. *Int. J. Cancer* 131, E33-44 (2012). WB, IP, Bovine, Human, Dog, Pig, Horse, Rabbit, Rat, Guinea pig, Mouse22072503
Ortega, M. C. et al. Megalin mediates the influence of sonic hedgehog on oligodendrocyte precursor cell migration and proliferation during development. *Glia* 60, 851-66 (2012). WB, IP, Bovine, Human, Dog, Pig, Horse, Rabbit, Rat, Guinea pig, Mouse22354480
Pandolfi, S. et al. WIP1 phosphatase modulates the Hedgehog signaling by enhancing GLI1 function. *Oncogene* 32, 4737-47 (2013). WB, IP, Bovine, Human, Dog, Pig, Horse, Rabbit, Rat, Guinea pig, Mouse23146903
Bosco-Clément, G. et al. Targeting Gli transcription activation by small molecule suppresses tumor growth. *Oncogene* 33, 2087-97 (2014). WB, IP, Bovine, Human, Dog, Pig, Horse, Rabbit, Rat, Guinea pig, Mouse23686308
Iwata, J. et al. Modulation of lipid metabolic defects rescues cleft palate in *Tgfr2* mutant mice. *Hum. Mol. Genet.* 23, 182-93 (2014). WB, IP, Bovine, Human, Dog, Pig, Horse, Rabbit, Rat, Guinea pig, Mouse23975680

Images




WB Suggested Anti-GLI1 Antibody Titration: 1.ug/ml
Positive Control: HepG2 cell lysate

Host:Rabbit
Target Name:GLI1



Host: Rabbit
Target Name: GLI1
Sample Tissue: Jurkat Cell Lysate
Antibody Dilution: 5.0µg/ml

Sample Tissue: Jurkat Whole Cell lysates
Antibody Dilution: 5.ug/ml



Host: Rabbit
Target Name: GLI1
Sample Tissue: HepG2 Cell Lysate
Antibody Dilution: 0.5µg/ml

Host: Rabbit
Target Name: GLI1
Sample Tissue: HepG2 Whole Cell lysates
Antibody Dilution: .5ug/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.