

# NR4A2 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI10410

#### **Product Information**

Application WB Primary Accession P43354

Other Accession <u>NM 006186, NP 006177</u>

**Reactivity** Human, Mouse, Rat, Zebrafish, Pig, Dog, Bovine

**Predicted** Human, Mouse, Rat, Zebrafish, Pig, Chicken, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 66591

#### **Additional Information**

**Gene ID** 4929

Alias Symbol HZF-3, NOT, NURR1, RNR1, TINUR

Other Names Nuclear receptor subfamily 4 group A member 2, Immediate-early response

protein NOT, Orphan nuclear receptor NURR1, Transcriptionally-inducible

nuclear receptor, NR4A2, NOT, NURR1, TINUR

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 50 ul of distilled water. Final anti-NR4A2 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

**Precautions** NR4A2 antibody - C-terminal region is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name NR4A2

Synonyms NOT, NURR1, TINUR

**Function** Transcriptional regulator which is important for the differentiation and

maintenance of meso-diencephalic dopaminergic (mdDA) neurons during development (PubMed: 15716272, PubMed: 17184956). It is crucial for expression of a set of genes such as SLC6A3, SLC18A2, TH and DRD2 which

are essential for development of mdDA neurons (By similarity).

**Cellular Location** Cytoplasm. Nucleus. Note=Mostly nuclear; oxidative stress promotes

cytoplasmic localization


**Tissue Location** 

Expressed in a number of cell lines of T-cell, B- cell and fibroblast origin. Strong expression in brain tissue

### References

Holla, V.R., (2006) J. Biol. Chem. 281 (5), 2676-2682 Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

## **Images**



WB Suggested Anti-NR4A2 Antibody Titration: .2-1 ug/ml Positive Control: Jurkat cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.