

ARID3A antibody - middle region

Rabbit Polyclonal Antibody Catalog # AI10430

Product Information

Application WB Primary Accession Q99856

Other Accession NM 005224, NP 005215

Reactivity Human, Mouse, Rat, Zebrafish, Pig, Dog, Bovine

Predicted Rat, Pig, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 62889

Additional Information

Gene ID 1820

Alias Symbol DRIL1, DRIL3, BRIGHT, E2FBP1

Other Names AT-rich interactive domain-containing protein 3A, ARID domain-containing

protein 3A, B-cell regulator of IgH transcription, Bright, Dead ringer-like protein 1, E2F-binding protein 1, ARID3A, DRIL1, DRIL3, DRX, E2FBP1

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 50 ul of distilled water. Final anti-ARID3A antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

Precautions ARID3A antibody - middle region is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name ARID3A

Synonyms DRIL1, DRIL3, DRX, E2FBP1

Function Transcription factor which may be involved in the control of cell cycle

progression by the RB1/E2F1 pathway and in B-cell differentiation.

Cellular Location Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00355,

ECO:0000269 | PubMed:17400556}. Cytoplasm Note=Shuttles between nucleus


and cytoplasm

Widely expressed, with highest expression in skeletal muscle, thalamus, and colon

References

Fukuyo, Y., et al., (2004) Cell Death Differ. 11 (7), 747-759Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images



WB Suggested Anti-ARID3A Antibody Titration: .2-1 ug/ml Positive Control: HepG2 cell lysate

ARID3A is supported by BioGPS gene expression data to be expressed in HepG2

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.