

# OVOL2 antibody - middle region

Rabbit Polyclonal Antibody Catalog # AI10479

#### **Product Information**

Application WB
Primary Accession Q9BRP0

Other Accession NM 021220, NP 067043

**Reactivity** Human, Mouse, Rat, Dog, Bovine

**Predicted** Human, Mouse, Rat, Pig, Chicken, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 30438

### **Additional Information**

**Gene ID** 58495

Alias Symbol EUROIMAGE566589, ZNF339

Other Names Transcription factor Ovo-like 2, hOvo2, Zinc finger protein 339, OVOL2,

**ZNF339** 

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 50 ul of distilled water. Final anti-OVOL2 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

**Precautions** OVOL2 antibody - middle region is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name OVOL2 ( HGNC:15804)

Synonyms ZNF339

**Function** Zinc-finger transcription repressor factor (PubMed: 19700410). Plays a critical

role in maintaining the identity of epithelial lineages by suppressing

epithelial-to mesenchymal transition (EMT) mainly through the repression of

ZEB1, an EMT inducer (By similarity). Positively regulates neuronal differentiation (By similarity). Suppresses cell cycling and terminal differentiation of keratinocytes by directly repressing MYC and NOTCH1 (PubMed: 19700410). Important for the correct development of primordial germ cells in embryos (By similarity). Plays dual functions in thermogenesis

and adipogenesis to maintain energy balance. Essential for brown/beige adipose tissue- mediated thermogenesis, is necessary for the development of brown adipocytes. In white adipose tissues, limits adipogenesis by blocking CEBPA binding to its transcriptional targets and inhibiting its transcription factor activity (By similarity).

**Cellular Location** Nucleus {ECO:0000250 | UniProtKB:Q8CIV7}.

**Tissue Location** Expressed in testis, ovary, heart and skeletal muscle (PubMed:12213202).


Expressed in the cornea, but absent from the corneal endothelium

(PubMed:26749309).

## References

Li,B., (2002) Genomics 80 (3), 319-325 Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

# **Images**



WB Suggested Anti-OVOL2 Antibody Titration: .2-1 ug/ml

ELISA Titer: 1:625

Positive Control: HT18 cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.