

Hmbox1 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI10533

Product Information

Application WB Primary Accession Q8BJA3

Other Accession <u>NM 177338</u>, <u>NP 796312</u>

Reactivity Human, Mouse, Rat, Zebrafish, Pig, Dog, Bovine

Predicted Human, Mouse, Rat, Zebrafish, Pig, Chicken, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 47116

Additional Information

Gene ID 219150

Alias Symbol AI451877, AI604847, F830020°C16Rik Other Names Homeobox-containing protein 1, Hmbox1

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 50 ul of distilled water. Final anti-Hmbox1 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

Precautions Hmbox1 antibody - C-terminal region is for research use only and not for use

in diagnostic or therapeutic procedures.

Protein Information

Name Hmbox1

Function Binds directly to 5'-TTAGGG-3' repeats in telomeric DNA (By similarity).

Associates with the telomerase complex at sites of active telomere processing and positively regulates telomere elongation (By similarity). Important for TERT binding to chromatin, indicating a role in recruitment of the telomerase complex to telomeres (PubMed: 23685356). Also plays a role in the alternative lengthening of telomeres (ALT) pathway in telomerase-negative cells where it promotes formation and/or maintenance of ALT-associated promyelocytic leukemia bodies (APBs) (By similarity). Enhances formation of telomere C-circles in ALT cells, suggesting a possible role in telomere recombination (By similarity). Might also be involved in the DNA damage response at telomeres

(By similarity).

Cellular Location

Nucleus. Cytoplasm {ECO:0000250 | UniProtKB:Q6NT76}. Chromosome, telomere. Nucleus, Cajal body {ECO:0000250 | UniProtKB:Q6NT76}. Nucleus, PML body {ECO:0000250 | UniProtKB:Q6NT76}. Note=Predominantly detected in cytoplasm. Localizes in a dynamic manner to actively processed telomeres. Localizes to the periphery of Cajal bodies. Associates with PML nuclear bodies in telomerase-negative cells {ECO:0000250 | UniProtKB:Q6NT76}

Images

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.