

KIF1A antibody - N-terminal region

Rabbit Polyclonal Antibody Catalog # AI10578

Product Information

WB
<u>Q12756</u>
<u>NM_004321</u> , <u>NP_004312</u>
Human, Mouse, Rat, Zebrafish, Pig, Dog, Bovine
Human, Mouse, Zebrafish, Pig, Chicken, Dog, Bovine
Rabbit
Polyclonal
191064

Additional Information

Gene ID	547
Alias Symbol	ATSV, C2orf20, DKFZp686I2094, FLJ30229, HUNC-104, MGC133285, MGC133286, UNC104, MRD9, HSN2C, SPG30
Other Names	Kinesin-like protein KIF1A, Axonal transporter of synaptic vesicles, Microtubule-based motor KIF1A, Unc-104- and KIF1A-related protein, hUnc-104, KIF1A, ATSV, C2orf20
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-KIF1A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	KIF1A antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	KIF1A
Synonyms	ATSV, C2orf20
Function	Kinesin motor with a plus-end-directed microtubule motor activity (By similarity). It is required for anterograde axonal transport of synaptic vesicle precursors (PubMed: <u>33880452</u>). Also required for neuronal dense core vesicles (DCVs) transport to the dendritic spines and axons. The interaction calcium-dependent with CALM1 increases vesicle motility and interaction with the scaffolding proteins PPFIA2 and TANC2 recruits DCVs to synaptic sites.

Cellular Location	Cytoplasm, cytoskeleton. Cell projection, neuron projection. Cell projection, axon {ECO:0000250 UniProtKB:P33173}. Cytoplasm, perinuclear region {ECO:0000250 UniProtKB:P33173}. Synapse {ECO:0000250 UniProtKB:P33173} Cytoplasmic vesicle, secretory vesicle, neuronal dense core vesicle membrane {ECO:0000250 UniProtKB:F1M4A4}; Peripheral membrane protein {ECO:0000250 UniProtKB:F1M4A4}; Cytoplasmic side {ECO:0000250 UniProtKB:F1M4A4}. Note=Within neuronal cells concentrated in the axon, with smaller amounts in the perinuclear and synaptic regions (By similarity). Accumulates at the distal tip of growing neurites. {ECO:0000250 UniProtKB:P33173, ECO:0000269 PubMed:25265257, ECO:0000269 PubMed:33880452}
Tissue Location	Expressed in neurons.

References

Kikkawa,M. (2006) EMBO J. 25 (18), 4187-4194Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images



WB Suggested Anti-KIF1A Antibody Titration: .2-1 ug/ml ELISA Titer: 1:625 Positive Control: THP-1 cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.