

# RNF13 antibody - N-terminal region

Rabbit Polyclonal Antibody Catalog # AI10611

#### **Product Information**

Application WB Primary Accession 043567

Other Accession NM 007282, NP 009213

**Reactivity**Human, Mouse, Rat, Pig, Dog, Bovine **Predicted**Human, Mouse, Rat, Chicken, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 42814

#### **Additional Information**

**Gene ID** 11342

Alias Symbol MGC13689, RZF

Other Names E3 ubiquitin-protein ligase RNF13, 6.3.2.-, RING finger protein 13, RNF13, RZF

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 50 ul of distilled water. Final anti-RNF13 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

**Precautions** RNF13 antibody - N-terminal region is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name RNF13 {ECO:0000303 | PubMed:18794910,

ECO:0000312 | HGNC:HGNC:10057}

**Function** E3 ubiquitin-protein ligase that regulates cell proliferation

(PubMed:<u>18794910</u>, PubMed:<u>23378536</u>, PubMed:<u>30595371</u>). Involved in apoptosis regulation (PubMed:<u>23378536</u>, PubMed:<u>30595371</u>). Mediates ER stress-induced activation of JNK signaling pathway and apoptosis by promoting ERN1 activation and splicing of XBP1 mRNA (PubMed:<u>23378536</u>, PubMed:<u>30595371</u>). Also involved in protein trafficking and localization

(PubMed:24387786).

**Cellular Location** Endoplasmic reticulum membrane; Single-pass type I membrane protein. Late

endosome membrane; Single-pass type I membrane protein. Lysosome

membrane; Single-pass type I membrane protein. Nucleus inner membrane {ECO:0000250 | UniProtKB:O54965}; Single-pass type I membrane protein. Note=Under certain conditions, relocalizes to recycling endosomes and to the inner nuclear membrane. {ECO:0000250 | UniProtKB:O54965}

#### **Tissue Location**

Widely expressed (at protein level). In normal pancreas, expressed in islets, but not in ducts, nor in acini (at protein level).

### References

Wesselkamper, S.C., (2006) Am. J. Respir. Cell Mol. Biol. 34 (1), 73-82 Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

## **Images**





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.