

HCFC2 antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI10654

Product Information

Application	WB
Primary Accession	Q9Y5Z7
Other Accession	NM_013320 , NP_037452
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	86779

Additional Information

Gene ID	29915
Alias Symbol	HCF-2, HCF2
Other Names	Host cell factor 2, HCF-2, C2 factor, HCFC2
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-HCFC2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	HCFC2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	HCFC2
Cellular Location	Cytoplasm. Nucleus
Tissue Location	Highly expressed in testis. Detected at lower levels in spleen, thymus, prostate, ovary, small intestine and colon

References

Yokoyama,A., (2004) Mol. Cell. Biol. 24 (13), 5639-5649 Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images



WB Suggested Anti-HCFC2 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Jurkat cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.