

# KCNH3 antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI10805

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">Q9ULD8</a>
<b>Other Accession</b>	<a href="#">NM_012284</a> , <a href="#">NP_036416</a>
<b>Reactivity</b>	Human, Mouse, Rat, Rabbit, Dog, Horse, Bovine
<b>Predicted</b>	Human, Rat, Pig, Horse, Bovine
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	117129

## Additional Information

---

<b>Gene ID</b>	23416
<b>Alias Symbol</b>	BEC1, ELK2, KIAA1282, Kv12.2
<b>Other Names</b>	Potassium voltage-gated channel subfamily H member 3, Brain-specific eag-like channel 1, BEC1, Ether-a-go-go-like potassium channel 2, ELK channel 2, ELK2, Voltage-gated potassium channel subunit Kv12.2, KCNH3, KIAA1282
<b>Format</b>	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
<b>Reconstitution &amp; Storage</b>	Add 50 ul of distilled water. Final anti-KCNH3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
<b>Precautions</b>	KCNH3 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information


---

<b>Name</b>	KCNH3 ( <a href="#">HGNC:6252</a> )
<b>Synonyms</b>	KIAA1282
<b>Function</b>	Pore-forming (alpha) subunit of a voltage-gated inwardly rectifying potassium channel (PubMed: <a href="#">10455180</a> ). Characterized by a fast rate of activation during depolarization followed by a rapid inactivation at much more depolarized value causing inward rectification due to a C-type inactivation mechanism (PubMed: <a href="#">10455180</a> ). Exhibits a rapid recovery from inactivation (PubMed: <a href="#">10455180</a> ).

<b>Cellular Location</b>	Cell membrane {ECO:0000250 UniProtKB:Q9WVJ0}; Multi-pass membrane protein {ECO:0000250 UniProtKB:Q9WVJ0} Note=Expression on the cell membrane requires at least one of the three glycosylation sites to carry a sugar chain irrespective of their positions. {ECO:0000250 UniProtKB:Q9WVJ0}
<b>Tissue Location</b>	Detected only in brain, in particular in the telencephalon (PubMed:10455180). Detected in the cerebral cortex, occipital pole, frontal and temporal lobe, putamen, amygdala, hippocampus and caudate nucleus (PubMed:10455180)

## Images

---



WB Suggested Anti-KCNH3 Antibody Titration: 0.2-1 µg/ml  
 ELISA Titer: 1:12500  
 Positive Control: OVCAR-3 cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.