

# KCNK10 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI10819

## **Product Information**

Application	WB
Primary Accession	<u>P57789</u>
Other Accession	<u>NM_021161, NP_066984</u>
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Horse, Bovine
Predicted	Human, Mouse, Rabbit, Pig, Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59765

## **Additional Information**

Gene ID	54207
Alias Symbol Other Names	TREK2, TREK-2, K2p10.1 Potassium channel subfamily K member 10, Outward rectifying potassium channel protein TREK-2, TREK-2 K(+) channel subunit, KCNK10, TREK2
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 100 ul of distilled water. Final anti-KCNK10 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	KCNK10 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

# Protein InformationNameKCNK10 {ECO:0000303 | PubMed:25766236,<br/>ECO:0000312 | HGNC:HGNC:6273}FunctionK(+) channel that conducts voltage-dependent outward rectifying currents<br/>upon membrane depolarization. Voltage sensing is coupled to K(+)<br/>electrochemical gradient in an 'ion flux gating' mode where outward but not<br/>inward ion flow opens the gate. Converts to voltage-independent 'leak'<br/>conductance mode upon stimulation by various stimuli including mechanical<br/>membrane stretch, acidic pH, heat and lipids (PubMed:10880510,<br/>PubMed:25766236, PubMed:26919430, PubMed:38605031). Homo- and<br/>heterodimerizes to form functional channels with distinct regulatory and<br/>gating properties (PubMed:30573346). In trigeminal ganglia sensory neurons,

	the heterodimer of KCNK10/TREK-2 and KCNK18/TRESK inhibits neuronal firing and neurogenic inflammation by stabilizing the resting membrane potential at K(+) equilibrium potential as well as by regulating the threshold of action potentials and the spike frequency (By similarity). Permeable to other monovalent ions such as Rb(+) and Cs(+) (PubMed: <u>26919430</u> ).
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:Q8BUW1}; Multi-pass membrane protein
Tissue Location	[Isoform A]: Abundantly expressed in pancreas and kidney and to a lower level in brain, testis, colon, and small intestine. In brain, mainly expressed in cerebellum, occipital lobe, putamen, and thalamus. No expression is detected in amygdala and spinal cord. [Isoform C]: Abundantly expressed in brain.

## References

Gu,W., et al., (2002) J. Physiol. (Lond.) 539 (PT 3), 657-668Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

### Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.