

KCNK10 antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI10819

Product Information

Application	WB
Primary Accession	P57789
Other Accession	NM_021161 , NP_066984
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Horse, Bovine
Predicted	Human, Mouse, Rabbit, Pig, Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59765

Additional Information

Gene ID	54207
Alias Symbol	TREK2, TREK-2, K2p10.1
Other Names	Potassium channel subfamily K member 10, Outward rectifying potassium channel protein TREK-2, TREK-2 K(+) channel subunit, KCNK10, TREK2
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 100 ul of distilled water. Final anti-KCNK10 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	KCNK10 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	KCNK10 {ECO:0000303 PubMed:25766236, ECO:0000312 HGNC:HGNC:6273}
Function	K(+) channel that conducts voltage-dependent outward rectifying currents upon membrane depolarization. Voltage sensing is coupled to K(+) electrochemical gradient in an 'ion flux gating' mode where outward but not inward ion flow opens the gate. Converts to voltage-independent 'leak' conductance mode upon stimulation by various stimuli including mechanical membrane stretch, acidic pH, heat and lipids (PubMed: 10880510 , PubMed: 25766236 , PubMed: 26919430 , PubMed: 38605031). Homo- and heterodimerizes to form functional channels with distinct regulatory and gating properties (PubMed: 30573346). In trigeminal ganglia sensory neurons,

the heterodimer of KCNK10/TREK-2 and KCNK18/TRESK inhibits neuronal firing and neurogenic inflammation by stabilizing the resting membrane potential at K(+) equilibrium potential as well as by regulating the threshold of action potentials and the spike frequency (By similarity). Permeable to other monovalent ions such as Rb(+) and Cs(+) (PubMed:[26919430](#)).

Cellular Location

Cell membrane {ECO:0000250 | UniProtKB:Q8BUW1}; Multi-pass membrane protein

Tissue Location

[Isoform A]: Abundantly expressed in pancreas and kidney and to a lower level in brain, testis, colon, and small intestine. In brain, mainly expressed in cerebellum, occipital lobe, putamen, and thalamus. No expression is detected in amygdala and spinal cord. [Isoform C]: Abundantly expressed in brain.

References

Gu,W., et al., (2002) J. Physiol. (Lond.) 539 (PT 3), 657-668
Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images



WB Suggested Anti-KCNK10 Antibody Titration: 2.5µg/ml
ELISA Titer: 1:62500
Positive Control: Jurkat cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.