10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

DDX19A antibody - N-terminal region

Rabbit Polyclonal Antibody
Catalog # AI10985

Product Information

Application	WB
Primary Accession	Q9NUU7
Other Accession	NM_018332, NP_060802
Reactivity	Human, Mouse, Rat, Dog, Horse
Predicted	Rat, Rabbit, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53975


Additional Information

Gene ID	55308
Alias Symbol	DDX19L, DDX19-DDX19L
Other Names	ATP-dependent RNA helicase DDX19A, 3.6.4.13, DDX19-like protein, DEAD box protein 19A, DDX19A, DDX19L
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 100 ul of distilled water. Final anti-DDX19A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	DDX19A antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DDX19A
Synonyms	DDX19L
Function	ATP-dependent RNA helicase involved in mRNA export from the nucleus. Rather than unwinding RNA duplexes, DDX19 functions as a remodeler of ribonucleoprotein particles, whereby proteins bound to nuclear mRNA are dissociated and replaced by cytoplasmic mRNA binding proteins.
Cellular Location	Cytoplasm {ECO:0000250|UniProtKB:Q9UMR2}. Nucleus, nucleoplasm {ECO:0000250|UniProtKB:Q9UMR2}. Note=Associates with the nuclear pore complex cytoplasmic fibrils {ECO:0000250|UniProtKB:Q9UMR2}

Images



WB Suggested Anti-DDX19A Antibody Titration: 2.5µg/ml
ELISA Titer: 1:62500
Positive Control: HepG2 cell lysate
DDX19A is strongly supported by BioGPS gene expression data to be expressed in Human HepG2 cells

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.